

# Inventor Search

Harris 09/871,491

=> d his

(FILE 'MEDLINE' ENTERED AT 14:00:22 ON 15 NOV 2001)  
DEL HIS Y

FILE 'MEDLINE, BIOSIS, WPIDS, HCAPLUS' ENTERED AT 14:03:30 ON 15 NOV 2001

E RAULET D/AU  
L1 305 S E3-4 OR E6-7  
E DIEFENBACH A/AU  
L2 61 S E3 OR E6-7  
L3 353 S L1 OR L2  
L4 139 S NKG2D OR NKG2 D  
L5 15 S L3 AND L4  
L6 7 DUP REM L5 (8 DUPLICATES REMOVED)

=> d bib ab 1-7

L6 ANSWER 1 OF 7 MEDLINE  
AN 2001468525 IN-PROCESS  
DN 21404050 PubMed ID: 11513138  
TI Strategies for target cell recognition by natural killer cells.  
AU **Diefenbach A; Raulet D H**  
CS Department of Molecular & Cell Biology, University of California Berkeley,  
94720-3200, USA.  
SO IMMUNOLOGICAL REVIEWS, (2001 Jun) 181 170-84.  
Journal code: GG4; 7702118. ISSN: 0105-2896.  
CY Denmark  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS IN-PROCESS; NONINDEXED; Priority Journals  
ED Entered STN: 20010830  
Last Updated on STN: 20010830  
AB Stimulation of natural killer (NK) cells is regulated by a complex balance  
of inhibitory and stimulatory receptors expressed by NK cells. However,  
the interaction of stimulatory receptors and their ligands is poorly  
understood. One stimulatory receptor, **NKG2D**, is expressed by all  
NK cells, stimulated CD8+ T cells, gammadelta T cells and macrophages.  
Recently, progress has been made in defining cellular ligands for  
**NKG2D**. Four different families of ligands have been identified in  
mice and humans, all of which are distantly related to MHC class I  
molecules. Some of the ligands are upregulated in transformed and infected  
cells, provoking an attack by the innate and adaptive immune systems. It  
appears that these "induced-self" ligands recognized by the **NKG2D**  
receptor may be a precedent for a new strategy of target cell recognition  
by the immune system.

L6 ANSWER 2 OF 7 MEDLINE DUPLICATE 1  
AN 2001510266 MEDLINE  
DN 21441910 PubMed ID: 11557981  
TI Rael and H60 ligands of the **NKG2D** receptor stimulate tumour  
immunity.  
AU **Diefenbach A; Jensen E R; Jamieson A M; Raulet D H**  
CS Department of Molecular and Cell Biology and Cancer Research Laboratory,  
University of California, Berkeley 94720, USA.  
SO NATURE, (2001 Sep 13) 413 (6852) 165-71.  
Journal code: NSC; 0410462. ISSN: 0028-0836.  
CY England; United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200110  
ED Entered STN: 20010917  
Last Updated on STN: 20011022

Entered Medline: 20011018

AB Natural killer (NK) cells attack many tumour cell lines, and are thought to have a critical role in anti-tumour immunity; however, the interaction between NK cells and tumour targets is poorly understood. The stimulatory lectin-like **NKG2D** receptor is expressed by NK cells, activated CD8+ T cells and by activated macrophages in mice. Several distinct cell-surface ligands that are related to class I major histocompatibility complex molecules have been identified, some of which are expressed at high levels by tumour cells but not by normal cells in adults. However, no direct evidence links the expression of these 'induced self' ligands with tumour cell rejection. Here we demonstrate that ectopic expression of the murine **NKG2D** ligands Raelbeta or H60 in several tumour cell lines results in potent rejection of the tumour cells by syngeneic mice. Rejection is mediated by NK cells and/or CD8+ T cells. The ligand-expressing tumour cells induce potent priming of cytotoxic T cells and sensitization of NK cells in vivo. Mice that are exposed to live or irradiated tumour cells expressing Rael or H60 are specifically immune to subsequent challenge with tumour cells that lack **NKG2D** ligands, suggesting application of the ligands in the design of tumour vaccines.

L6 ANSWER 3 OF 7 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:210647 BIOSIS

DN PREV200100210647

TI The ligands for mouse **NKG2D** are expressed on tumor cells and activate NK cells and macrophages.

AU Diefenbach, A. (1); Jamieson, A. M. (1); Raulet, D. H. (1)

CS (1) Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California Berkeley, Berkeley, CA USA

SO Immunobiology, (November, 2000) Vol. 203, No. 1-2, pp. 436. print.

Meeting Info.: Joint Annual Meeting of the German and Dutch Societies of Immunology Dusseldorf, Germany November 29-December 02, 2000  
ISSN: 0171-2985.

DT Conference

LA English

SL English

L6 ANSWER 4 OF 7 MEDLINE

DUPLICATE 2

AN 2001216034 MEDLINE

DN 21205390 PubMed ID: 11248803

TI Ligands for the murine **NKG2D** receptor: expression by tumor cells and activation of NK cells and macrophages.

CM Comment in: Nat Immunol. 2000 Aug;1(2):95-7

AU Diefenbach A; Jamieson A M; Liu S D; Shastri N; Raulet D H

CS Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California, Berkeley, USA.

SO Nat Immunol, (2000 Aug) 1 (2) 119-26.

Journal code: DOG; 100941354. ISSN: 1529-2908.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200105

ED Entered STN: 20010521

Last Updated on STN: 20010521

Entered Medline: 20010517

AB Natural killer (NK) cells attack tumor and infected cells, but the receptors and ligands that stimulate them are poorly understood. Here we report the expression cloning of two murine ligands for the lectin-like receptor **NKG2D**. The two ligands, H-60 and Rael beta, are distant relatives of major histocompatibility complex class I molecules. **NKG2D** ligands are not expressed by most normal cells but are

up-regulated on numerous tumor cells. We show that mouse **NKG2D** is expressed by NK cells, activated CD8+ T cells and activated macrophages. Expression of either **NKG2D** ligand by target cells triggers NK cell cytotoxicity and interferon-gamma secretion by NK cells, as well as nitric oxide release and tumor necrosis factor alpha transcription by macrophages. Thus, through their interaction with **NKG2D**, H-60 and Rael beta are newly identified potent stimulators of innate immunity.

L6 ANSWER 5 OF 7 MEDLINE DUPLICATE 3  
 AN 2000045048 MEDLINE  
 DN 20045048 PubMed ID: 10574749  
 TI Natural killer cells: stress out, turn on, tune in.  
 AU **Diefenbach A; Raulet D H**  
 CS Department of Molecular and Cell Biology, Cancer Research Laboratory, 485 Life Sciences Addition, University of California at Berkeley, Berkeley, 94720-3200, USA.  
 SO CURRENT BIOLOGY, (1999 Nov 18) 9 (22) R851-3. Ref: 14  
 Journal code: B44; 9107782. ISSN: 0960-9822.  
 CY ENGLAND: United Kingdom  
 DT Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LA English  
 FS Priority Journals  
 EM 200005  
 ED Entered STN: 20000518  
 Last Updated on STN: 20000518  
 Entered Medline: 20000511  
 AB Natural killer cells attack tumor cells, infected cells and some normal cells, but the basis of their specificity is not completely understood. Recent studies indicate that epithelial tumor cells upregulate a stress-induced MHC class-I-like protein termed MICA, triggering NK cells via a recently described receptor called **NKG2D**.

L6 ANSWER 6 OF 7 BIOSIS COPYRIGHT 2001 BIOSIS  
 AN 1999:219315 BIOSIS  
 DN PREV199900219315  
 TI CD94 and NKG2 lectin-like receptors on mouse natural killer cells.  
 AU Vance, Russell E. (1); Kraft, Jennifer; Altman, John; Jensen, Peter;  
**Raulet, David H. (1)**  
 CS (1) University of California (Berkeley), Berkeley, CA USA  
 SO Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 91.  
 Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity  
 Seventeenth International Natural Killer Cell Workshop Warrenton,  
 Virginia, USA October 17-21, 1998  
 ISSN: 1018-8916.  
 DT Conference  
 LA English

L6 ANSWER 7 OF 7 MEDLINE DUPLICATE 4  
 AN 1998124458 MEDLINE  
 DN 98124458 PubMed ID: 9464811  
 TI Cloning of a mouse homolog of CD94 extends the family of C-type lectins on murine natural killer cells.  
 AU Vance R E; Tanamachi D M; Hanke T; **Raulet D H**  
 CS Department of Molecular and Cell Biology & Cancer Research Laboratory, University of California at Berkeley, 94720, USA.  
 NC R01 AI35021 (NIAID)  
 SO EUROPEAN JOURNAL OF IMMUNOLOGY, (1997 Dec) 27 (12) 3236-41.  
 Journal code: EN5; 1273201. ISSN: 0014-2980.  
 CY GERMANY: Germany, Federal Republic of  
 DT Journal; Article; (JOURNAL ARTICLE)

LA English  
 FS Priority Journals  
 OS GENBANK-AF030311; GENBANK-AF030312; GENBANK-AF030313  
 EM 199802  
 ED Entered STN: 19980306  
 Last Updated on STN: 19980306  
 Entered Medline: 19980220

AB Two families of major histocompatibility complex (MHC) class I-specific receptors are found on natural killer (NK) cells: immunoglobulin-like receptors and C-type lectin receptors. In mice, the latter category is represented by the Ly49 family of receptors, whereas in humans, NK cells express the distantly related CD94, which forms MHC class I-specific heterodimers with NKG2 family members. Humans also express the MHC class I-specific p50/p58/p70 family of immunoglobulin-like receptors, but these have not been identified in mice. Hence, there is no known instance of an MHC class I-specific receptor that is expressed by both human and murine NK cells. Here we report the cloning of CD94 from the CB.17 and C57BL/6 strains of mice. Mouse CD94 is 54% identical and 66% similar to human CD94, and is also a member of the C-type lectin superfamily. Mouse CD94 is expressed efficiently on the cell surface of cells transiently transfected with the corresponding cDNA, but surface CD94 was unable to mediate detectable binding to MHC class I-expressing ConA blasts. Notably, mouse CD94, like human CD94, has a very short cytoplasmic tail, suggesting the existence of partner chains that may play a role in ligand binding and signaling. Like many other C-type lectins expressed by NK cells, mouse CD94 maps to the NK complex on distal chromosome 6, synteneic to human CD94. We also demonstrate that mouse CD94 is highly expressed specifically by mouse NK cells, raising the possibility that mice, like humans, express multiple families of MHC class I-specific receptors on their NK cells. Murine homologs of human NKG2 family members have not yet been identified, but we report here the existence of a murine **NKG2D**-like sequence that also maps to the murine NK complex near CD94 and Ly49 family members.

=> d his

(FILE 'HOME' ENTERED AT 13:44:42 ON 15 NOV 2001)

FILE 'HCAPLUS' ENTERED AT 13:44:46 ON 15 NOV 2001

L1 27 S NKG2D  
L2 0 S NK G2D  
L3 45 S NKG2D/AB  
L4 81 S NKG2  
L5 8 S L4 (2W) D  
L6 8 S (NKG2 D)/AB  
L7 57 S L1 OR L3 OR L5 OR L6  
L8 360944 S TUMOR OR CARCINOMA? OR CANCER# OR NEOPLAS? OR MELENOMA?  
L9 367140 S L8 OR MELANOMA?  
L10 9 S L7 AND L8  
L11 9 S L7 AND L9  
L12 111394 S LIGAND#  
L13 9 S L12 AND L7  
L14 7 S L13 NOT L11

FILE 'WPIDS' ENTERED AT 13:51:09 ON 15 NOV 2001

L15 1 S NKG2D OR NKG2 D

FILE 'BIOSIS' ENTERED AT 13:51:37 ON 15 NOV 2001

L16 58 S NKG2D OR NKG2 D  
L17 1015943 S TUMOR OR CARCINOMA? OR CANCER# OR NEOPLAS? OR MELANOMA?  
L18 16 S L16 AND L17  
L19 25 S LIGAND# AND L16  
L20 15 S L19 NOT L18

FILE 'HCAPLUS, BIOSIS' ENTERED AT 13:53:05 ON 15 NOV 2001

L21 21 DUP REM L11 L18 (4 DUPLICATES REMOVED)  
L22 18 DUP REM L14 L20 (4 DUPLICATES REMOVED)

=&gt; d.bib ab 121 1-21;d bib ab 122 1-18

L21 ANSWER 1 OF 21 HCAPLUS COPYRIGHT 2001 ACS

AN 2001:713586 HCAPLUS

DN 135:269070

TI Multifunctional proteins binding to **NKG2D** receptor complex and their use in treatment of **cancer**, infections, and autoimmune diseases

IN Kufer, Peter; Riethmueller, Gert; Lutterbuese, Ralf; Borschert, Katrin; Kischel, Roman; Mayer, Monika; Hofmeister, Robert

PA Germany

SO PCT Int. Appl., 114 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001071005	A2	20010927	WO 2001-EP3414	20010326
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				

PRAI EP 2000-106467 A 20000324

AB The present invention relates to a multifunctional polypeptide comprising a first domain comprising a binding site specifically recognizing an extracellular epitope of the **NKG2D** receptor complex and a second domain having receptor or ligand function. Furthermore, the present invention relates to polynucleotides encoding the multifunctional polypeptide, to vectors comprising said polypeptides and to cells comprising said polynucleotides or said vectors. The invention also relates to compns. comprising either of the above recited mols., alone or in combination, as well as to specific medical uses of the multifunctional polypeptide of the invention. Thus, scFv proteins binding to **NKG2D** and Ep-CAM are produced. These scFv's recruited cytotoxic lymphocytes (CD8+ T cells and NK cells) and caused lysis of Ep-CAM-producing cells.

L21 ANSWER 2 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:385890 BIOSIS

DN PREV200100385890

TI Receptors involved in human NK cell activation in the process of natural cytotoxicity.

AU Moretta, Lorenzo (1); Biassoni, Roberto; Bottino, Cristina; Mingari, Maria Cristina; Moretta, Alessandro

CS (1) Istituto Giannina Gaslini, 16148, Genova Italy

SO Cooper, Max D.; Takai, Toshiyuki; Ravetch, Jeffrey V.. (2001) pp. 199-209.

Activating and inhibitory immunoglobulin-like receptors. print.

Publisher: Springer-Verlag Tokyo Inc. 3-13 Hongo 3-chome, Bunkyo-ku, Tokyo, 113-0033, Japan.

Meeting Info.: CREST International Symposium on Immunoglobulin-like Receptors Sendai City, Japan September 18-19, 2000

ISBN: 4-431-70297-0 (cloth).

DT Book; Conference

LA English

SL English

L21 ANSWER 3 OF 21 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 1  
 AN 2001:729002 HCAPLUS  
 TI Ectopic expression of retinoic acid early inducible-1 gene (RAE-1) permits natural killer cell-mediated rejection of a MHC class I-bearing tumor in vivo  
 AU Cerwenka, Adelheid; Baron, Jody L.; Lanier, Lewis L.  
 CS Department of Microbiology and Immunology and the Cancer Research Institute, University of California, San Francisco, CA, 94143-0414, USA  
 SO Proc. Natl. Acad. Sci. U. S. A. (2001), 98(20), 11521-11526  
 CODEN: PNASA6; ISSN: 0027-8424  
 PB National Academy of Sciences  
 DT Journal  
 LA English  
 AB In 1986, Karre and colleagues reported that natural killer (NK) cells rejected an MHC class I-deficient tumor cell line (RMA-S) but they did not reject the same cell line if it expressed MHC class I (RMA). Based on this observation, they proposed the concept that NK cells provide immune surveillance for "missing self," e.g., they eliminate cells that have lost class I MHC antigens. This seminal observation predicted the existence of inhibitory NK cell receptors for MHC class I. Here, we present evidence that NK cells are able to reject tumors expressing MHC class I if the tumor expresses a ligand for **NKG2D**. Mock-transfected RMA cells resulted in tumor formation. In contrast, when RMA cells were transfected with the retinoic acid early inducible gene-1 .gamma. or .delta. (RAE-1), ligands for the activating receptor **NKG2D**, the tumors were rejected. The tumor rejection was mediated by NK cells, and not by CD1-restricted NK1.1+ T cells. No T cell-mediated immunol. memory against the parental tumor was generated in the animals that had rejected the RAE-1 transfected tumors, which succumbed to rechallenge with the parental RMA tumor. Therefore, NK cells are able to reject a tumor expressing RAE-1 mols., despite expression of self MHC class I on the tumor, demonstrating the potential for NK cells to participate in immunity against class I-bearing malignancies.

RE.CNT 33

RE

- (1) Bahram, S; Adv Immunol 2000, V76, P1 HCAPLUS
  - (2) Bakker, A; Hum Immunol 2000, V61, P18 HCAPLUS
  - (3) Bauer, S; Science 1999, V285, P727 HCAPLUS
  - (4) Biron, C; Annu Rev Immunol 1999, V17, P189 HCAPLUS
  - (5) Bix, M; Nature (London) 1991, V349, P329 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 4 OF 21 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 2  
 AN 2001:298002 HCAPLUS  
 DN 135:60134  
 TI Role of **NKG2D** in tumor cell lysis mediated by human NK cells: cooperation with natural cytotoxicity receptors and capability of recognizing tumors of nonepithelial origin  
 AU Pende, Daniela; Cantoni, Claudia; Rivera, Paola; Vitale, Massimo; Castriconi, Roberta; Marcenaro, Stefania; Nanni, Marina; Biassoni, Roberto; Bottino, Cristina; Moretta, Alessandro; Moretta, Lorenzo  
 CS Istituto Nazionale per la Ricerca sul Cancro, Genoa, Italy  
 SO Eur. J. Immunol. (2001), 31(4), 1076-1086  
 CODEN: EJIMAF; ISSN: 0014-2980  
 PB Wiley-VCH Verlag GmbH  
 DT Journal  
 LA English  
 AB **NKG2D** is a recently described activating receptor expressed by both NK cells and CTL. In this study we investigated the role of **NKG2D** in the natural cytotoxicity mediated by NK cell clones. The role of **NKG2D** varied depending on the type of target cells analyzed. Lysis of various tumors appeared to be exclusively natural cytotoxicity receptors (NCR) dependent. In contrast, killing of another

group of target cells, including not only the epithelial cell lines HELA and IGROV-1, but also the FO-1 melanoma, the JA3 leukemia, the Daudi Burkitt lymphoma and even normal PHA-induced lymphoblasts, involved both NCR and **NKG2D**. Notably, NK cell clones expressing low surface densities of NCR(NCRdull) could lyse these tumors in an exclusively **NKG2D**-dependent fashion. Remarkably, not all of these targets expressed MICA/B, thus implying the existence of addnl. ligands recognized by **NKG2D**, possibly represented by GPI-linked mols. Finally, we show that the engagement of different HLA class I-specific inhibitory receptors by either specific antibodies or the appropriate HLA class I ligand led to inhibition of **NKG2D**-mediated NK cell triggering.

RE.CNT 37

RE

- (1) Bauer, S; Science 1999, V285, P727 HCAPLUS
  - (2) Borrego, F; J Exp Med 1998, V187, P813 HCAPLUS
  - (3) Braud, V; Nature 1998, V391, P795 HCAPLUS
  - (4) Cantoni, C; Eur J Immunol 1998, V28, P327 HCAPLUS
  - (5) Cantoni, C; J Exp Med 1999, V189, P787 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 5 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:267825 BIOSIS

DN PREV200100267825

TI Expression of **NKG2D** and MDL-1 on porcine myeloid cells.

AU Yim, Daesong (1); Jie, Hyun-Bae (1); Sotiriadis, John (1); Kim, Yoon-Sang (1); Shin, Soon Cheon (1); Lanier, Lewis L.; Kim, Yoon B. (1)

CS (1) Finch University of Health Sciences/The Chicago Medical School, 3333 Green Bay Road, North Chicago, IL, 60064 USA

SO FASEB Journal, (March 7, 2001) Vol. 15, No. 4, pp. A692. print.  
 Meeting Info.: Annual Meeting of the Federation of American Societies for Experimental Biology on Experimental Biology 2001 Orlando, Florida, USA  
 March 31-April 04, 2001  
 ISSN: 0892-6638.

DT Conference

LA English

SL English

AB **NKG2D** is a C-type lectin family receptor that associates with DAP10, recognizing tumor antigens such as MICA/B, RAE-1 and H60. We cloned porcine **NKG2D** receptor cDNA by RT-PCR. Porcine **NKG2D** cDNA has an open reading frame of 642 bp. Its expected polypeptide sequence is 214 amino acids. Porcine **NKG2D** has 66% sequence identity with human **NKG2D** and 56% identity with mouse **NKG2D**. RT-PCR analysis reveals that porcine **NKG2D** transcripts are expressed in PBL, NK cells, macrophages, and monocytes, but not in granulocytes. LPS upregulated **NKG2D** mRNA expression in macrophages. Porcine **NKG2D** gene is located on chromosome 5q25. When transiently transfected into COS-7 cells, porcine **NKG2D** requires DAP10 for cell surface expression. Myeloid DAP12-associating lectin-1 (MDL-1) is a type II membrane protein that associates with DAP12. Two isoforms of porcine MDL-1 cDNA were cloned from pulmonary alveolar macrophages. Porcine MDL-1 short form has 165 amino acids and 70% sequence identity with mouse MDL-1 short form. The long form has 20 more amino acids in the stalk region and 71% sequence identity with human MDL-1 and 67% with mouse MDL-1 long form. Porcine MDL-1 contains a conserved lysine in the transmembrane domain. MDL-1 transcripts were detected exclusively in macrophages and monocytes by RT-PCR. When transfected into 293 cells, porcine MDL-1 is expressed on the cell surface associated with DAP12. MDL-1 mRNA was detected in PBMC of neonatal "germ-free" piglets. Thus, MDL-1 may be involved in innate immunity.

L21 ANSWER 6 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:275629 BIOSIS

DN PREV200100275629



- TI **NKG2D/DAP10: An immune receptor for oncofetal antigens.**  
 AU Cerwenka, Adelheid (1); Lanier, Lewis L. (1)  
 CS (1) UCSF, San Francisco, CA, 94143 USA  
 SO FASEB Journal, (March 7, 2001) Vol. 15, No. 4, pp. A660. print.  
 Meeting Info.: Annual Meeting of the Federation of American Societies for  
 Experimental Biology on Experimental Biology 2001 Orlando, Florida, USA  
 March 31-April 04, 2001  
 ISSN: 0892-6638.
- DT Conference  
 LA English  
 SL English  
 AB NK cells are important effector cells in innate immunity providing protection against certain viral infections and tumors. A delicate balance between activating and inhibitory signals dictates their recognition of abnormal, distressed cells. We identified a family of cell surface proteins encoded by the retinoic acid inducible (RAE-1) genes and the H60 minor histocompatibility antigen that function as ligands for the mouse activating receptor **NKG2D**. Expression of these ligands is low in healthy adult tissue, but high in the embryo and on several **tumor** cell lines. Cell surface expression of RAE-1 molecules, which have low homology to classical MHC class 1 molecules, does not require the TAP transporter or beta2-microglobulin. Ectopic expression of RAE-1 in a MHC class 1 positive cell line confers target susceptibility to NK cell attack. Thus, the interaction between the receptor **NKG2D** expressed on NK cells with its ligands delivers a strong activating signal, which is able to overcome the self-MHC class 1 mediated inhibitory signals. These data emphasize the importance of the **NKG2D**-ligand interaction in NK cell activation and suggest a potential role in anti-**tumor** immunity.
- L21 ANSWER 7 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS  
 AN 2001:536998 BIOSIS  
 DN PREV200100536998  
 TI Regulation of cutaneous malignancy by gammadelta T cells.  
 AU Girardi, Michael; Oppenheim, David E.; Steele, Carrie R.; Lewis, Julia M.; Glusac, Earl; Filler, Renata; Hobby, Paul; Sutton, Brian; Tigelaar, Robert E.; Hayday, Adrian C. (1)  
 CS (1) Peter Gorer Department of Immunobiology, Guy's King's St. Thomas' Medical School, King's College, London, SE1 9RT: adrian.hayday@kcl.ac.uk UK  
 SO Science (Washington D C), (19 October, 2001) Vol. 294, No. 5542, pp. 605-609. print.  
 ISSN: 0036-8075.
- DT Article  
 LA English  
 SL English  
 AB The localization of gammadelta T cells within epithelia suggests that these cells may contribute to the down-regulation of epithelial malignancies. We report that mice lacking gammadelta cells are highly susceptible to multiple regimens of cutaneous carcinogenesis. After exposure to carcinogens, skin cells expressed Rae-1 and H60, major histocompatibility complex-related molecules structurally resembling human MICA. Each of these is a ligand for **NKG2d**, a receptor expressed by cytolytic T cells and natural killer (NK) cells. In vitro, skin-associated **NKG2d**+ gammadelta cells killed skin **carcinoma** cells by a mechanism that was sensitive to blocking **NKG2d** engagement. Thus, local T cells may use evolutionarily conserved proteins to negatively regulate malignancy.
- L21 ANSWER 8 OF 21 HCAPLUS COPYRIGHT 2001 ACS  
 AN 2001:296377 HCAPLUS  
 DN 135:75404  
 TI Activating receptors and coreceptors involved in human natural killer

- cell-mediated cytotoxicity
- AU Moretta, Alessandro; Bottino, Cristina; Vitale, Massimo; Pende, Daniela; Cantoni, Claudia; Mingari, Maria Cristina; Biassoni, Roberto; Moretta, Lorenzo
- CS Dip. Med. Sper., Universita degli Studi di Genova, Genoa, Italy
- SO< Annu. Rev. Immunol. (2001), 19, 197-223
- CODEN: ARIMDU; ISSN: 0732-0582
- PB Annual Reviews Inc.
- DT Journal; General Review
- LA English
- AB A review and discussion with 127 refs. Natural killer cells can discriminate between normal cells and cells that do not express adequate amts. of major histocompatibility complex (MHC) class I mols. The discovery, both in mouse and in human, of MHC-specific inhibitory receptors clarified the mol. basis of this important NK cell function. However, the triggering receptors responsible for pos. NK cell stimulation remained elusive until recently. Some of these receptors have now been identified in humans, thus shedding some light on the mol. mechanisms involved in NK cell activation during the process of natural cytotoxicity. Three novel, NK-specific, triggering surface mols. (NKp46, NKp30, and NKp44) have been identified. They represent the first members of a novel emerging group of receptors collectively termed natural cytotoxicity receptors (NCR). A direct correlation exists between the surface d. of NCR and the ability of NK cells to kill various tumors. NKp46 is the only NCR involved in human NK-mediated killing of murine target cells. Accordingly, a homolog of NKp46 has been detected in mouse. Mol. cloning of NCR revealed novel members of the Ig superfamily displaying a low degree of similarity to each other and to known human mols. NCRs are coupled to different signal transducing adaptor proteins, including CD3.zeta., Fc.epsilon.RI.gamma., and KARAP/DAP12. Another triggering NK receptor is **NKG2D**. It appears to play either a complementary or a synergistic role with NCRs. Thus, the triggering of NK cells in the process of tumor cell lysis may often depend on the concerted action of NCR and **NKG2D**. In some instances, however, it may uniquely depend upon the activity of NCR or **NKG2D** only. Strict **NKG2D**-dependency can be appreciated using clones that, in spite of their NCRdull phenotype, efficiently lyse certain epithelial tumors or leukemic cell lines. Other triggering surface mols. including 2B4 and the novel NKp80 appear to function as coreceptors rather than as true receptors. Indeed, they can induce natural cytotoxicity only when co-engaged with a triggering receptor. While an altered expression or function of NCR or **NKG2D** is being explored as a possible cause of immunol. disorders, 2B4 dysfunction has already been assocd. with a severe form of immunodeficiency. Indeed, in patients with the X-linked lymphoproliferative disease, the inability to control Epstein-Barr virus infections may be consequent to a major dysfunction of 2B4 that exerts inhibitory instead of activating functions.

RE.CNT 111

RE

- (1) Auchincloss, H; Annu Rev Immunol 1998, V16, P433 HCAPLUS
  - (2) Bakker, A; Immunity 2000, V13, P345 HCAPLUS
  - (3) Biassoni, R; Eur J Immunol 1997, V27, P3095 HCAPLUS
  - (4) Biassoni, R; Eur J Immunol 1999, V29, P1014 HCAPLUS
  - (5) Biassoni, R; J Exp Med 1996, V183, P645 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 9 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:482112 BIOSIS

DN PREV200100482112

TI Rael and H60 ligands of the **NKG2D** receptor stimulate tumour immunity.

AU Diefenbach, Andreas; Jensen, Eric R.; Jamieson, Amanda M.; Raulet, David H. (1)

CS (1) Department of Molecular and Cell Biology and Cancer Research  
Laboratory, University of California, 485 Life Sciences Addition,  
Berkeley, CA, 94720: raulet@uclink4.berkeley.edu USA

SO Nature (London), (13 September, 2001) Vol. 413, No. 6852, pp. 165-171.  
print.  
ISSN: 0028-0836.

DT Article

LA English

SL English

AB Natural killer (NK) cells attack many tumour cell lines, and are thought  
to have a critical role in anti-tumour immunity; however, the interaction  
between NK cells and tumour targets is poorly understood. The stimulatory  
lectin-like **NKG2D** receptor is expressed by NK cells, activated  
CD8+ T cells and by activated macrophages in mice. Several distinct  
cell-surface ligands that are related to class I major histocompatibility  
complex molecules have been identified, some of which are expressed at  
high levels by tumour cells but not by normal cells in adults. However, no  
direct evidence links the expression of these 'induced self' ligands with  
tumour cell rejection. Here we demonstrate that ectopic expression of the  
murine **NKG2D** ligands Raelbeta or H60 in several tumour cell  
lines results in potent rejection of the tumour cells by syngeneic mice.  
rejection is mediated by NK cells and/or CD8+ T cells. The  
ligand-expressing tumour cells induce potent priming of cytotoxic T cells  
and sensitization of NK cells in vivo. Mice that are exposed to live or  
irradiated tumour cells expressing Rael or H60 are specifically immune to  
subsequent challenge with tumour cells that lack **NKG2D** ligands,  
suggesting application of the ligands in the design of tumour vaccines.

L21 ANSWER 10 OF 21 HCAPLUS COPYRIGHT 2001 ACS      DUPLICATE 3

AN 2001:229621 HCAPLUS

DN 135:18196

TI Recognition of **tumor** cells by the innate immune system

AU Soloski, Mark J.

CS Division of Rheumatology and the Program in Immunology, Department of  
Medicine, Johns Hopkins University School of Medicine, Baltimore, MD,  
21205, USA

SO Curr. Opin. Immunol. (2001), 13(2), 154-162  
CODEN: COPIEL; ISSN: 0952-7915

PB Elsevier Science Ltd.

DT Journal; General Review

LA English

AB A review with 88 refs. There has been a rapid increase in the authors'  
understanding of the cellular components of the innate immune system, the  
receptors used to distinguish changes in homeostasis, and how these  
components integrate into an anti-tumor effector response. Recently,  
significant progress has been made in the identification of ligands for  
receptors that activate NK cells, and the results have implications for  
the recognition of tumor cells.

RE.CNT 88

RE

- (1) Aldrich, C; Cell 1994, V79, P649 HCAPLUS
  - (2) Amadou, C; Immunol Rev 1999, V167, P211 HCAPLUS
  - (3) Bakker, A; Immunity 2000, V13, P345 HCAPLUS
  - (4) Bakker, A; Proc Natl Acad Sci USA 1999, V96, P9792 HCAPLUS
  - (5) Bauer, S; Science 1999, V285, P727 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 11 OF 21 HCAPLUS COPYRIGHT 2001 ACS

AN 2001:297982 HCAPLUS

DN 135:59877

TI ULBPs, novel MHC class I-related molecules, bind to CMV glycoprotein UL16  
and stimulate NK cytotoxicity through the **NKG2D** receptor

AU Cosman, David; Mullberg, Jurgen; Sutherland, Claire L.; Chin, Wilson;

CS Armitage, Richard; Fanslow, William; Kubin, Marek; Chalupny, N. Jan  
 Department of Molecular Biology, Immunex Corporation, Seattle, WA, 98101,  
 USA  
 SO Immunity (2001), 14(2), 123-133  
 CODEN: IUNIEH; ISSN: 1074-7613  
 PB Cell Press  
 DT Journal  
 LA English  
 AB The human cytomegalovirus glycoprotein, UL16, binds to two members of a  
 novel family of mols., the ULBPs, and to the MHC class I homolog, MICB.  
 The ULBPs are GPI-linked glycoproteins belonging to the extended MHC class  
 I family but are only distantly related to MICB. The ULBP and MICB mols.  
 are ligands for the activating receptor, **NKG2D**/DAP10, and this  
 interaction is blocked by a sol. form of UL16. The ULBPs stimulate  
 cytokine and chemokine prodn. from NK cells, and expression of ULBPs in NK  
 cell-resistant target cells confers susceptibility to NK cell  
 cytotoxicity. Masking of NK cell recognition of ULBP or MIC antigens by  
 UL16 provides a potential mechanism by which human cytomegalovirus-  
 infected cells might evade attack by the immune system.

RE.CNT 45

RE

- (1) Ando, H; Immunogenetics 1997, V46, P499 HCAPLUS
  - (2) Bahram, S; Proc Natl Acad Sci USA 1994, V91, P6259 HCAPLUS
  - (3) Bauer, S; Science 1999, V285, P727 HCAPLUS
  - (4) Baum, P; EMBO J 1994, V13, P3992 HCAPLUS
  - (5) Biron, C; Annu Rev Immunol 1999, V17, P189 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 12 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS  
 AN 2001:37271 BIOSIS  
 DN PREV200100037271  
 TI Soluble forms of the novel, MHC class 1-related molecules, ULBP1 and  
 ULBP2, bind to, and functionally activate NK cells.  
 AU Chalupny, J. (1); Cosman, D. (1); Mullberg, J. (1); Chin, W. (1);  
 Cassiano, L. (1); Means, G. (1); Derry, J. (1); Russell, C. (1); Armitage,  
 R. (1); Sutherland, C. (1); Fanslow, W. (1); Kubin, M. (1)  
 CS (1) Immunex Corp, Seattle, WA, 98101 USA  
 SO FASEB Journal, (April 20, 2000) Vol. 14, No. 6, pp. A1018. print.  
 Meeting Info.: Joint Annual Meeting of the American Association of  
 Immunologists and the Clinical Immunology Society Seattle, Washington, USA  
 May 12-16, 2000  
 ISSN: 0892-6638.  
 DT Conference  
 LA English  
 SL English

L21 ANSWER 13 OF 21 HCAPLUS COPYRIGHT 2001 ACS  
 AN 2000:552552 HCAPLUS  
 TI Exposing tumor cells to killer cell attack  
 AU Watzl, C.; Long, E. O.  
 CS Laboratory of Immunogenetics National Institute of Allergy and Infectious  
 Diseases, National Institutes of Health, Rockville, MD, 20852, USA  
 SO Nat. Med. (N. Y.) (2000), 6(8), 867-868  
 CODEN: NAMEFI; ISSN: 1078-8956  
 PB Nature America Inc.  
 DT Journal  
 LA English  
 AB Natural killer (NK) cells attack tumor and virally infected cells in the  
 absence of antigen presentation, utilizing a combination of signals from  
 activation and inhibitory receptors. Recent reports have identified the  
 NK and T-cell surface protein **NKG2D** as a receptor for tumor cell  
 ligands that activates killing of tumor targets.

RE.CNT 9

RE

- (1) Bauer, S; Science 1999, V285, P727 HCAPLUS
- (2) Cerwenka, A; Immunity 2000, V12, P721 HCAPLUS
- (3) Chang, C; J Immunol 1999, V163, P4651 HCAPLUS
- (4) Groh, V; Proc Natl Acad Sci 1999, V96, P6879 HCAPLUS
- (5) Ljunggren, H; Immunol Today 1990, V11, P237 HCAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 14 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:210647 BIOSIS

DN PREV200100210647

TI The ligands for mouse **NKG2D** are expressed on tumor cells and activate NK cells and macrophages.

AU Diefenbach, A. (1); Jamieson, A. M. (1); Raulet, D. H. (1)

CS (1) Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California Berkeley, Berkeley, CA USA

SO Immunobiology, (November, 2000) Vol. 203, No. 1-2, pp. 436. print.  
Meeting Info.: Joint Annual Meeting of the German and Dutch Societies of Immunology Dusseldorf, Germany November 29-December 02, 2000  
ISSN: 0171-2985.

DT Conference

LA English

SL English

L21 ANSWER 15 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2000:238053 BIOSIS

DN PREV200000238053

TI A single amino acid substitution causes loss of expression of a MICA allele.

AU Li, Zihai; Groh, Veronika; Strong, Roland K.; Spies, Thomas (1)

CS (1) Fred Hutchinson Cancer Research Center, 1100 Fairview Ave. N., Seattle, WA, 98109 USA

SO Immunogenetics, (March, 2000) Vol. 51, No. 3, pp. 246-248.  
ISSN: 0093-7711.

DT Article

LA English

SL English

L21 ANSWER 16 OF 21 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 4

AN 2000:557581 HCAPLUS

DN 133:236795

TI Ligands for the murine **NKG2D** receptor: expression by tumor cells and activation of NK cells and macrophages

AU Diefenbach, Andreas; Jamieson, Amanda M.; Liu, Scot D.; Shastri, Nilabh; Raulet, David H.

CS Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California, Berkeley, CA, USA

SO Nat. Immunol. (2000), 1(2), 119-126  
CODEN: NIAMCZ; ISSN: 1529-2908

PB Nature America Inc.

DT Journal

LA English

AB Natural killer (NK) cells attack tumor and infected cells, but the receptors and ligands that stimulate them are poorly understood. Here we report the expression cloning of two murine ligands for the lectin-like receptor **NKG2D**. The two ligands, H-60 and Rae-1.beta., are distant relatives of major histocompatibility complex class I mols. **NKG2D** ligands are not expressed by most normal cells but are up-regulated on numerous tumor cells. We show that mouse **NKG2D** is expressed by NK cells, activated CD8+T cells and activated macrophages. Expression of either **NKG2D** ligand by target cells triggers NK cell cytotoxicity and interferon-.gamma. secretion by NK cells, as well as nitric oxide release and tumor necrosis factor .alpha. transcription by

macrophages. Thus, through their interaction with **NKG2D**, H-60 and Rael.beta. are newly identified potent stimulators of innate immunity.

RE.CNT 41

RE

- (1) Altman, J; Science 1996, V274, P94 HCAPLUS
  - (2) Amadou, C; Immunol Rev 1999, V167, P211 HCAPLUS
  - (3) Bahram, S; Proc Natl Acad Sci USA 1994, V91, P6259 HCAPLUS
  - (4) Bauer, S; Science 1999, V285, P727 HCAPLUS
  - (5) Brown, M; J Exp Med 1998, V188, P2083 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 17 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2000:28283 BIOSIS

DN PREV200000028283

TI Natural killer cells: Stress out, turn on, tune in.

AU Diefenbach, Andreas (1); Raulet, David H. (1)

CS (1) Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California at Berkeley, 485 Life Sciences Addition, Berkeley, CA, 94720-3200 USA

SO Current Biology, (Nov. 18, 1999) Vol. 9, No. 22, pp. R851-R853.  
ISSN: 0960-9822.

DT Article

LA English

SL English

AB Natural killer cells attack **tumor** cells, infected cells and some normal cells, but the basis of their specificity is not completely understood. Recent studies indicate that epithelial **tumor** cells upregulate a stress-induced MHC class-I-like protein termed MICA, triggering NK cells via a recently described receptor called **NKG2D**

L21 ANSWER 18 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1999:484525 BIOSIS

DN PREV199900484525

TI An activating immunoreceptor complex formed by **NKG2D** and DAP10.

AU Wu, Jun; Song, Yaoli; Bakker, Alexander B.H.; Bauer, Stefan; Spies, Thomas; Lanier, Lewis L. (1); Phillips, Joseph H.

CS (1) DNAX Research Institute, 901 California Avenue, Palo Alto, CA, 94304 USA

SO Science (Washington D C), (July 30, 1999) Vol. 285, No. 5428, pp. 730-732.  
ISSN: 0036-8075.

DT Article

LA English

SL English

AB Many immune receptors are composed of separate ligand-binding and signal-transducing subunits. In natural killer (NK) and T cells, DAP10 was identified as a cell surface adaptor protein in an activating receptor complex with **NKG2D**, a receptor for the stress-inducible and **tumor**-associated major histocompatibility complex molecule MICA. Within the DAP10 cytoplasmic domain, an Src homology 2 (SH2) domain-binding site was capable of recruiting the p85 subunit of the phosphatidylinositol 3-kinase (PI 3-kinase), providing for **NKG2D**-dependent signal transduction. Thus, **NKG2D**-DAP10 receptor complexes may activate NK and T cell responses against MICA-bearing tumors.

L21 ANSWER 19 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1999:484524 BIOSIS

DN PREV199900484524

TI Activation of NK cells and T cells by **NKG2D**, a receptor for stress-inducible MICA.

AU Bauer, Stefan; Groh, Veronika; Wu, Jun; Steinle, Alexander; Phillips, Joseph H.; Lanier, Lewis L.; Spies, Thomas (1)

CS (1) Clinical Research Division, Fred Hutchinson Cancer Research Center,  
1100 Fairview Avenue North, Seattle, WA, 98109 USA  
SO Science (Washington D C), (July 30, 1999) Vol. 285, No. 5428, pp. 727-729.  
ISSN: 0036-8075.

DT Article

LA English

SL English

AB Stress-inducible MICA, a distant homolog of major histocompatibility complex (MHC) class I, functions as an antigen for gammadelta T cells and is frequently expressed in epithelial tumors. A receptor for MICA was detected on most gammadelta T cells, CD8+ alphabeta T cells, and natural killer (NK) cells and was identified as **NKG2D**. Effector cells from all these subsets could be stimulated by ligation of **NKG2D**. Engagement of **NKG2D** activated cytolytic responses of gammadelta T cells and NK cells against transfectants and epithelial tumor cells expressing MICA. These results define an activating immunoreceptor-MHC ligand interaction that may promote antitumor NK and T cell responses.

L21 ANSWER 20 OF 21 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1996:316201 BIOSIS

DN PREV199699038557

TI An autosomal dominant locus, Nka, mapping to the Ly-49 region of a rat natural killer (NK) gene complex, controls NK cell lysis of allogeneic lymphocytes.

AU Dissen, Erik (1); Ryan, James C.; Seaman, William E.; Fossum, Sigbjorn  
CS (1) Dep. Anat., Inst. Basic Medical Sciences, Univ. Oslo, P.O. Box 1105, Blindern, N-0137 Oslo Norway

SO Journal of Experimental Medicine, (1996) Vol. 183, No. 5, pp. 2197-2207.  
ISSN: 0022-1007.

DT Article

LA English

AB Natural Killer (NK) cells can recognize and kill MHC-incompatible normal bone marrow-derived cells. Presently characterized MHC-binding receptors on NK cells, including the Ly-49 family in the mouse, transmit inhibitory signals upon binding to cognate class I MHC ligands. Here we study in vivo NK-mediated lysis of normal allogeneic lymphocytes in crosses between alloreactivity-competent PVG rats and alloreactivity-deficient DA rats. NK cells from both strains are able to lyse standard tumor targets. We identify an autosomal dominant locus, Nka, that controls NK-mediated alloreactivity. Individuals carrying the dominant PVG allele in single dose were fully competent in eliminating allogeneic target cells, suggesting that Nka encodes or regulates a gene product inducing or activating alloreactivity. By linkage analysis and pulsed field gel electrophoresis, a natural killer gene complex (NKC) on rat chromosome 4 is described that contains the rat NKR-P1 and Ly-49 multigene families plus a rat **NKG2D** homologue. Nka maps within the NKC, together with the most telomeric Ly-49 family members, but separate from **NKG2D** and the NKR-P1 family. The Nka-encoded response, moreover, correlates with the expression of transcripts for Ly-49 receptors in NK cell populations, as Northern blot analysis demonstrated low expression of Ly-49 genes in DA NK cells, in contrast to high expression in alloreactivity-competent PVG, (DA times PVG)F-1, and PVG.1AV1 NK cells. The low Ly-49 expression in DA is not induced by MHC haplotype, as demonstrated by high expression of Ly-49 in the DA MHC-congenic PVG.1AV1 strain. Finally, we have cloned and characterized the first four members of the rat Ly-49 gene family. Their cytoplasmic domains demonstrate substantial heterogeneity, consistent with the hypothesis that different Ly-49 family members may subserve different signaling functions.

L21 ANSWER 21 OF 21 HCAPLUS COPYRIGHT 2001 ACS

AN 1993:162459 HCAPLUS

DN 118:162459

TI NKG2 proteins of natural killer cells, cDNA encoding them, and methods for treatment of **cancer** or virus infection  
 IN Houchins, Jeffrey P.; Yabe, Toshio; McSherry, Cynthia M.; Bach, Fritz H.; Hofer, Erhard  
 PA University of Minnesota, USA; Sandoz Ltd.  
 SO PCT Int. Appl., 61 pp.  
 CODEN: PIXXD2

DT Patent  
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9217198	A1	19921015	WO 1992-US2469	19920327
	W: JP, US				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE				
	EP 585257	A1	19940309	EP 1992-909331	19920327
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	JP 06506358	T2	19940721	JP 1992-508930	19920327
	US 6262244	B1	20010717	US 1995-543246	19951013
PRAI	US 1991-676663	A2	19910328		
	WO 1992-US2469	W	19920327		
	US 1993-122514	B1	19930924		

AB The cDNAs for 4 NKG2 proteins of human natural killer cells are cloned and sequenced. Antibodies to these proteins; chimeric antibodies recognizing NKG2 protein and a cancer- or virus-specific antigen; NKG2-cytotoxic protein fusion proteins; and the use of these antibodies or chimeric proteins for treatment of cancer or virus infection are claimed. The 4 cDNAs represent a new mammalian gene family. These proteins displayed significant sequence similarity only with type II transmembrane proteins with C-type animal lectin domains. The NKG2-C protein was manufd. with transgenic Sf9 cells and the extracellular domain of this protein manufd. as a fusion protein in Escherichia coli.

L22 ANSWER 1 OF 18 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 1

AN 2001:330267 HCAPLUS

DN 135:106137

TI Complex structure of the activating immunoreceptor **NKG2D** and its MHC class I-like **ligand** MICA

AU Li, Pingwei; Morris, Daniel L.; Willcox, Benjamin E.; Steinle, Alexander; Spies, Thomas; Strong, Roland K.

CS Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, 98109, USA

SO Nat. Immunol. (2001), 2(5), 443-451

CODEN: NIAMCZ; ISSN: 1529-2908

PB Nature America Inc.

DT Journal

LA English

AB The major histocompatibility complex (MHC) class I homolog, MICA, is a stress-inducible ligand for **NKG2D**, a C-type lectin-like activating immunoreceptor. The crystal structure of this ligand-receptor complex that the authors report here reveals an **NKG2D** homodimer bound to a MICA monomer in an interaction that is analogous to that seen in T cell receptor-MHC class I protein complexes. Similar surfaces on each **NKG2D** monomer interact with different surfaces on either the .alpha.1 or .alpha.2 domains of MICA. The binding interactions are large in area and highly complementary. The central section of the .alpha.2-domain helix, disordered in the structure of MICA alone, is ordered in the complex and forms part of the **NKG2D** interface. The extensive flexibility of the interdomain linker of MICA is shown by its altered conformation when crystd. alone or in complex with



**NKG2D.**

RE.CNT 53

RE

- (1) Bahram, S; Immunogenetics 1996, V43, P230 HCAPLUS
  - (2) Bahram, S; Proc Natl Acad Sci 1994, V91, P6259 HCAPLUS
  - (4) Bauer, S; Science 1999, V285, P727 HCAPLUS
  - (5) Berman, H; Nucleic Acids Res 2000, V28, P235 HCAPLUS
  - (6) Boyington, J; Immunity 1999, V10, P75 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 2 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:385993 BIOSIS

DN PREV200100385993

TI MICA and MICB genes: Can the enigma of their polymorphism be resolved.

AU Stephens, Henry A. F. (1)

CS (1) Institute of Urology and Nephrology, University College London, Middlesex Hospital, 48 Riding House Street, London, W1W 7EY: h.stephens@ucl.ac.uk UK

SO Trends in Immunology, (July, 2001) Vol. 22, No. 7, pp. 378-385. print. ISSN: 1471-4906.

DT General Review

LA English

SL English

AB The human MHC class I chain-related genes (MICA and MICB) are located within the HLA class I region of chromosome 6. Their organization, expression and products differ considerably from classical HLA class I genes. MIC proteins are considered to be markers of 'stress' in the epithelia, and act as **ligands** for cells expressing a common activatory natural killer-cell receptor (**NKG2D**). Molecular models are now available for the MICA protein, both bound and complexed with **NKG2D**. MICA molecules appear to be highly flexible and polymorphic, although the functional relevance and implications of their polymorphism have yet to be fully discerned.

L22 ANSWER 3 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:291228 BIOSIS

DN PREV200100291228

TI Crystal structure of the complex between the stress-inducible MHC class I homolog MIC-A and the NK cell receptor **NKG2D**.

AU Strong, Roland K. (1); Li, Pingwei (1); Morris, Daniel L. (1); Steinle, Alexander (1); Spies, Thomas (1)

CS (1) Fred Hutchinson Cancer Research Center, Basic Sciences, 1100 Fairview Ave. North, Seattle, WA, 98109 USA

SO FASEB Journal, (March 7, 2001) Vol. 15, No. 4, pp. A321. print. Meeting Info.: Annual Meeting of the Federation of American Societies for Experimental Biology on Experimental Biology 2001 Orlando, Florida, USA March 31-April 04, 2001 ISSN: 0892-6638.

DT Conference

LA English

SL English

AB The major histocompatibility complex (MHC) class I homolog MIC-A functions as a stress-inducible antigen, expressed on intestinal epithelium and epithelially-derived tumors, that is broadly recognized by the Vdelta1-bearing subset of gammadelta T cells, CD8+ alphabeta T cells and NK cells independent of beta2-microglobulin and bound peptides. MIC-A recognition by these cells is mediated through interactions with the stimulatory NK receptor **NKG2D**, a divergent member of the C-type lectin-like family of proteins and a distant relative of other members of the NKG2 family of NK cell receptors and CD94. The crystal structure of the complex reveals that an **NKG2D** homodimer binds to a MIC-A monomer, contacting residues of both the alpha1 and alpha2 domains, in an interaction distinct from other NK receptor/**ligand** complex

structures. A section of the alpha2 domain helix, disordered in the structure of MIC-A crystallized on its own, is ordered in the complex and forms part of the MIC-A/NKG2D interface. Potential receptor binding sites on the underside of the platform, on the side opposite the surface recognized by alphabeta T cell receptors on MHC class I/peptide complexes, proposed on the basis of the MIC-A crystal structure, do not form part of the MIC-A/NKG2D interface in the crystal structure of the complex. Rather, these sites participate in extensive MIC-A/MIC-A crystal contacts yielding a MIC-A tetramer that recapitulates the interdomain relationship seen in conventional, beta2-microglobulin-binding MHC class I homologs.

L22 ANSWER 4 OF 18 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 2

AN 2001:481688 HCAPLUS

TI Interactions of human NKG2D with its ligands MICA, MICB, and homologs of the mouse RAE-1 protein family

AU Steinle, Alexander; Li, Pingwei; Morris, Daniel L.; Groh, Veronika; Lanier, Lewis L.; Strong, Roland K.; Spies, Thomas

CS Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, WA, 98109, USA

SO Immunogenetics (2001), 53(4), 279-287  
CODEN: IMNGBK; ISSN: 0093-7711

PB Springer-Verlag

DT Journal

LA English

AB NKG2D is an activating receptor that is expressed on most natural killer (NK) cells, CD8 .alpha..beta. T cells, and .gamma..delta. T cells. Among its ligands is the distant major histocompatibility complex class I homolog MICA, which has no function in antigen presentation but is induced by cellular stress. To extend previous functional evidence, the NKG2D-MICA interaction was studied in isolation. NKG2D homodimers formed stable complexes with monomeric MICA in soln., demonstrating that no other components were required to facilitate this interaction. MICA glycosylation was not essential but enhanced complex formation. Sol. NKG2D also bound to cell surface MICB, which has structural and functional properties similar to those of MICA. Moreover, NKG2D stably interacted with surface mols. encoded by three newly identified cDNA sequences (N2DL-1, -2, and -3), which are identical to the human ULBP proteins and may represent homologs of the mouse retinoic acid-early inducible family of NKG2D ligands. Because of the substantial sequence divergence among these mols., these results indicated promiscuous modes of receptor binding. Comparison of allelic variants of MICA revealed large differences in NKG2D binding that were assocd. with a single amino acid substitution at position 129 in the .alpha.2 domain. Varying affinities of MICA alleles for NKG2D may affect thresholds of NK-cell triggering and T-cell modulation.

RE.CNT 41

RE

(1) Bahram, S; Immunogenetics 1996, V43, P230 HCAPLUS

(2) Bahram, S; Proc Natl Acad Sci 1994, V91, P6259 HCAPLUS

(4) Bauer, S; Science 1999, V285, P727 HCAPLUS

(5) Bjorkman, P; Annu Rev Biochem 1990, V59, P253 HCAPLUS

(6) Bjorkman, P; Nature 1987, V329, P506 HCAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 5 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:410147 BIOSIS

DN PREV200100410147

TI Costimulation of CD8alphabeta T cells by NKG2D via engagement by MIC induced on virus-infected cells.

AU Groh, Veronika (1); Rhinehart, Rebecca; Randolph-Habecker, Julie; Topp, Max S.; Riddell, Stanley R.; Spies, Thomas (1)

CS (1) Clinical Research Division, Fred Hutchinson Cancer Research Center,  
1100 Fairview Ave. North, Seattle, WA, 98109: vgroh@fred.fhcrc.org,  
tspies@fred.fhcrc.org USA  
SO Nature Immunology, (March, 2001) Vol. 2, No. 3, pp. 255-260. print.  
ISSN: 1529-2908.  
DT Article  
LA English  
SL English  
AB **NKG2D** is an activating receptor that stimulates innate immune  
responses by natural killer cells upon engagement by MIC **ligands**  
, which are induced by cellular stress. Because **NKG2D** is also  
present on most CD8alphabeta T cells, it may modulate antigen-specific T  
cell responses, depending on whether MIC molecules-distant homologs of  
major histocompatibility complex (MHC) class I with no function in antigen  
presentation-are induced on the surface of pathogen-infected cells. We  
found that infection by cytomegalovirus (CMV) resulted in substantial  
increases in MIC on cultured fibroblast and endothelial cells and was  
associated with induced MIC expression in interstitial pneumonia. MIC  
engagement of **NKG2D** potentially augmented T cell antigen receptor  
(TCR)-dependent cytolytic and cytokine responses by CMV-specific CD28-  
CD8alphabeta T cells. This function overcame viral interference with MHC  
class I antigen presentation. Combined triggering of TCR-CD3 complexes and  
**NKG2D** induced interleukin 2 production and T cell proliferation.  
Thus **NKG2D** functioned as a costimulatory receptor that can  
substitute for CD28.

L22 ANSWER 6 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:410146 BIOSIS

DN PREV200100410146

TI Crystal structure of the murine NK cell-activating receptor **NKG2D**  
at 1.95 Å.

AU Wolan, Dennis W.; Teyton, Luc; Rudolph, Markus G.; Villmow, Brigitte;  
Bauer, Stefan; Busch, Dirk H.; Wilson, Ian A. (1)

CS (1) Department of Molecular Biology, Skaggs Institute for Chemical  
Biology, Scripps Research Institute, 10550 North Torrey Pines Road, La  
Jolla, CA, 92037: dirk.busch@lrz.tum.de, wilson@scripps.edu USA

SO Nature Immunology, (March, 2001). Vol. 2, No. 3, pp. 248-254. print.  
ISSN: 1529-2908.

DT Article

LA English

SL English

AB **NKG2D**, a homodimeric lectin-like receptor, is a unique  
stimulatory molecule that is found on natural killer cells, T cells and  
activated macrophages. The natural **ligands** for murine  
**NKG2D** are distant major histocompatibility complex homologs,  
retinoic acid early transcript (Rae1) and H-60 minor histocompatibility  
antigen. The crystal structure of the extracellular region of murine  
**NKG2D** reveals close homology with other C-type lectin receptors  
such as CD94, Ly49A, rat MBP-A and CD69. However, the precise mode of  
dimeric assembly varies among these natural killer receptors, as well as  
their surface topography and electrostatic properties. The **NKG2D**  
structure provides the first structural insights into the role and  
**ligand** specificity of this stimulatory receptor in the innate and  
adaptive immune system.

L22 ANSWER 7 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:454125 BIOSIS

DN PREV200100454125

TI Molecular competition for **NKG2D**: H60 and RAE1 compete unequally  
for **NKG2D** with dominance of H60.

AU O'Callaghan, Christopher A.; Cerwenka, Adelheid; Willcox, Benjamin E.;  
Lanier, Lewis L.; Bjorkman, Pamela J. (1)

CS (1) Howard Hughes Medical Institute, California Institute of Technology,

- Pasadena, CA, 91125: bjorkman@its.caltech.edu USA
- SO Immunity, (August, 2001) Vol. 15, No. 2, pp. 201-211. print.  
ISSN: 1074-7613.
- DT Article
- LA English
- SL English
- AB **NKG2D** is a potent activating receptor on natural killer cells, T cells, and macrophages. Mouse **NKG2D** interacts with two cell surface ligands related to class I MHC molecules: RAE1 and H60. We used soluble versions of **NKG2D**, RAE1, and H60 to characterize their interactions. RAE1 and H60 each bind **NKG2D** with nanomolar affinities, indicating tighter binding than most cell surface immune interactions, but **NKG2D** binds to H60 with approx25-fold higher affinity than to RAE1. RAE1 and H60 compete directly for occupancy of **NKG2D**, and, thus, **NKG2D** can be occupied by only one ligand at a time. The **NKG2D**-H60 interaction is more temperature dependent and makes greater use of electrostatic interactions than the **NKG2D**-RAE1 interaction. The distinct thermodynamic profiles provide insights into the different molecular mechanisms of the binding interactions.
- L22 ANSWER 8 OF 18 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 3
- AN 2001:659192 HCAPLUS
- TI The UL16-binding proteins, a novel family of MHC class I-related ligands for **NKG2D**, activate natural killer cell functions
- AU Sutherland, Claire L.; Chalupny, N. Jan; Cosman, David
- CS Department of Molecular Biology, Immunex Corporation, Seattle, WA, 98101, USA
- SO Immunol. Rev. (2001), 181, 185-192  
CODEN: IMRED2; ISSN: 0105-2896
- PB Munksgaard International Publishers Ltd.
- DT Journal
- LA English
- AB The UL16-binding proteins (ULBPs) are a novel family of MHC class I-related mols. (MICs) that were identified based on their ability to bind to the human cytomegalovirus (HCMV) glycoprotein UL16. UL16 also binds to a member of another family of MHC class I-like mols., MICB. The ULBPs and MICs are ligands for **NKG2D**/DAP10, an activating receptor expressed by natural killer (NK) cells and other immune effector cells, and this interaction can be blocked by UL16. Engagement of **NKG2D**/DAP10 by ULBPs or MICs expressed on a target cell can overcome an inhibitory signal generated by NK-cell recognition of MHC class I mols. and trigger NK cytotoxicity. ULBPs elicit their effects on NK cells by activating the janus kinase 2, signal transducer and activator of transcription 5, extracellular-signal-regulated kinase mitogen-activated protein kinase and Akt/protein kinase B signal transduction pathways. Although ULBPs alone activate multiple signaling pathways and induce modest cytokine prodn., ULBPs synergize strongly with interleukin-12 for prodn. of interferon- $\gamma$  by NK cells. This finding is consistent with reports in T cells that **NKG2D**/DAP10 can act as a costimulatory receptor in a similar manner as CD28. The possible roles of ULBPs in mediating immune responses to viruses and tumors and the potential mechanisms by which UL16 may allow HCMV to evade immune detection are areas of active investigation.
- RE.CNT 50
- RE
- (1) Azuma, M; J Immunol 1993, V150, P1147 HCAPLUS
- (2) Bauer, S; Science 1999, V285, P727 HCAPLUS
- (3) Biron, C; Annu Rev Immunol 1999, V17, P189 HCAPLUS
- (5) Blery, M; Hum Immunol 2000, V61, P51 HCAPLUS
- (6) Boise, L; Immunity 1995, V3, P87 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 9 OF 18 HCAPLUS COPYRIGHT 2001 ACS

AN 2001:707947 HCAPLUS

TI Rael and H60 ligands of the **NKG2D** receptor stimulate tumour immunity

AU Diefenbach, Andreas; Jansen, Eric R.; Jamison, Amanda M.; Raulet, David H.

CS Department of Molecular and Cell Biology and Cancer Research Lab.,  
University of California, Berkeley, 94720, USA

SO Nature (London, U. K.) (2001), 413(6852), 165-171  
CODEN: NATUAS; ISSN: 0028-0836

PB Nature Publishing Group

DT Journal

LA English

AB Natural killer (NK) cells attack many tumor cell lines, and are thought to have a crit. role in anti-tumor immunity; however, the interaction between NK cells and tumor targets is poorly understood. The stimulatory lectin-like **NKG2D** receptor is expressed by NK cells, activated CD8+ T cells and by activated macrophages in mice. Several distinct cell-surface ligands that are related to class I major histocompatibility complex mols. have been identified, some of which are expressed at high levels by tumor cells but not by normal cells in adults. However, no direct evidence links the expression of these 'induced self' ligands with tumor cell rejection. Here we demonstrate that ectopic expression of the murine **NKG2D** ligands Rael.beta. or H60 in several tumor cell lines results in potent rejection of the tumor cells by syngeneic mice. Rejection is mediated by NK cells and/or CD8+ T cells. The ligand-expressing tumor cells induce potent priming of cytotoxic T cells and sensitization of NK cells in vivo. Mice that are exposed to live or irradiated tumor cells expressing Rael or H60 are specifically immune to subsequent challenge with tumor cells that lack **NKG2D** ligands, suggesting application of the ligands in the design of tumor vaccines.

RE.CNT 30

RE

(1) Bauer, S; Science 1999, V285, P727 HCAPLUS

(2) Cerwenka, A; Immunity 2000, V12, P721 HCAPLUS

(3) Cosman, D; Immunity 2001, V14, P123 HCAPLUS

(4) Diefenbach, A; Nature Immunol 2000, V1, P119 HCAPLUS

(5) Dranoff, G; Proc Natl Acad Sci 1993, V90, P3539 HCAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 10 OF 18 HCAPLUS COPYRIGHT 2001 ACS

AN 2001:659190 HCAPLUS

TI Ligands for natural killer cell receptors: Redundancy or specificity

AU Cerwenka, Adelheid; Lanier, Lewis L.

CS Department of Microbiology and Immunology and the Cancer Research  
Institute, University of California, San Francisco, CA, 94143-0414, USA

SO Immunol. Rev. (2001), 181, 158-169  
CODEN: IMRED2; ISSN: 0105-2896

PB Munksgaard International Publishers Ltd.

DT Journal

LA English

AB Several inhibitory and activating receptors involved in natural killer cell activation have been characterized. The increasing knowledge about their ligands, including classical MHC class I mols., non-classical MHC class I mols. and MHC class I-related mols., is shedding new light on the targets of innate immune recognition. While classical MHC class I mols. are constitutively expressed, some MHC class I-related (MIC) mols., however, are stress-induced by ill-defined stimuli. Two families of ligands for the human activating **NKG2D** receptor have been identified. These are the MIC proteins encoded by two highly polymorphic genes within the MHC class I and the retinoic acid-inducible early gene-1-like (also designated UL16-binding) proteins encoded by genes

outside the MHC. For the mouse **NKG2D** receptor, one family, contg. at least five distinct ligands, has been described. A better understanding about how targets signal their distress, which renders them susceptible to natural killer (NK)-cell attack, will help to define the role of NK cells in antimicrobial and antitumor immunity and transplantation.

RE.CNT 90

RE

- (1) Bahram, S; Adv Immunol 2000, V76, P1 HCAPLUS
  - (2) Bahram, S; Proc Natl Acad Sci USA 1994, V91, P6259 HCAPLUS
  - (3) Bakker, A; Hum Immunol 2000, V61, P18 HCAPLUS
  - (4) Bakker, A; Immunity 2000, V13, P345 HCAPLUS
  - (5) Bauer, S; Science 1999, V285, P727 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 11 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:141877 BIOSIS

DN PREV200100141877

TI ULBPs, Novel MHC class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the **NKG2D** receptor.

AU Cosman, David (1); Mullberg, Jurgen; Sutherland, Claire L.; Chin, Wilson; Armitage, Richard; Fanslow, William; Kubin, Marek; Chalupny, N. Jan

CS (1) Department of Molecular Biology, Immunex Corporation, 51 University Street, Seattle, WA, 98101: cosman@immunex.com USA

SO Immunity, (February, 2001) Vol. 14, No. 2, pp. 123-133. print.  
ISSN: 1074-7613.

DT Article

LA English

SL English

AB The human cytomegalovirus glycoprotein, UL16, binds to two members of a novel family of molecules, the ULBPs, and to the MHC class I homolog, MICB. The ULBPs are GPI-linked glycoproteins belonging to the extended MHC class I family but are only distantly related to MICB. The ULBP and MICB molecules are **ligands** for the activating receptor, **NKG2D** /DAPI0, and this interaction is blocked by a soluble form of UL16. The ULBPs stimulate cytokine and chemokine production from NK cells, and expression of ULBPs in NK cell-resistant target cells confers susceptibility to NK cell cytotoxicity. Masking of NK cell recognition of ULBP or MIC antigens by UL16 provides a potential mechanism by which human cytomegalovirus-infected cells might evade attack by the immune system.

L22 ANSWER 12 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2001:106283 BIOSIS

DN PREV200100106283

TI Triggering receptors involved in natural killer cell-mediated cytotoxicity against choriocarcinoma cell lines.

AU Sivori, Simona; Parolini, Silvia; Marcenaro, Emanuela; Millo, Romano; Bottino, Cristina; Moretta, Alessandro (1)

CS (1) Dipartimento di Medicina Sperimentale, Sezione di Istologia, Universita di Genova, Via G.B. Marsano 10, 16132, Genova: alemoret@unige.it/bottino@ermes.cba.unige.it Italy

SO Human Immunology, (November, 2000) Vol. 61, No. 11, pp. 1055-1058. print.  
ISSN: 0198-8859.

DT Article

LA English

SL English

AB The lack of classical HLA-class I molecules on trophoblast is necessary to prevent allorecognition by maternal CTL, but may induce activation of NK cells. A protective role against NK cells equipped of suitable inhibitory receptors has been proposed for nonclassical HLA-class I molecules including HLA-E and HLA-G. In the present study we show that the NK-mediated killing of two choriocarcinoma cell lines, JAR and JEG3, is induced upon engagement of natural cytotoxicity receptors (NCR) with their

specific **ligands**. In particular, we show that NKp44, a triggering receptor expressed at the NK cell surface only after in vitro culture in the presence of IL-2, plays a central role in triggering NK cytotoxicity against trophoblast cells. Also NKp46 appear to contribute to this function by cooperating with NKp44. On the other hand, other triggering receptors such as NKp30, 2B4, and **NKG2D** are not involved in killing of choriocarcinoma. Our findings suggest that resistance of trophoblast to NK-mediated cytotoxicity is the result of insufficient activating interactions between the various triggering NK receptors and their target cell **ligands**. On the other hand, the interaction of nonclassical HLA class I molecules with inhibitory NK receptors appears to play only a marginal role in regulating the susceptibility of choriocarcinoma to NK mediated cytotoxicity.

L22 ANSWER 13 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS  
 AN 2001:385618 BIOSIS  
 DN PREV200100385618  
 TI Human natural killer cell activating receptors.  
 AU Biassoni, Roberto (1); Cantoni, Claudia; Falco, Michela; Pende, Daniela; Millo, Romano; Moretta, Lorenzo; Bottino, Cristina; Moretta, Alessandro  
 CS (1) Istituto Nazionale per la Ricerca sul Cancro, Laboratorio di Immunologia, IST/CBA, L.go R. Benzi, 10, 16132, Genova: biassoni@cba.unige.it Italy  
 SO Molecular Immunology, (December, 2000) Vol. 37, No. 17, pp. 1015-1024.. print.  
 ISSN: 0161-5890.  
 DT General Review  
 LA English  
 SL English  
 AB Natural killer (NK) cells were poorly characterized until 10 years ago and few molecules expressed on their cell surface were known. Now the situation has changed dramatically, since a plethora of receptors characterized by opposite functions have been functionally and molecularly defined. NK cells express clonally distributed inhibitory receptors specific for different groups of HLA class I alleles, thus protecting normal cells from NK-mediated lysis. On the contrary, various activating receptors are involved in triggering of NK-mediated natural cytotoxicity. Their engagement induces human NK cells to kill target cells that are either HLA class I-negative or -deficient. Here a brief description of the activating receptors and coreceptor and of their **ligand(s)** is given.

L22 ANSWER 14 OF 18 HCAPLUS COPYRIGHT 2001 ACS DUPLICATE 4  
 AN 2000:472382 HCAPLUS  
 DN 133:191846  
 TI Retinoic acid early inducible genes define a **ligand** family for the activating **NKG2D** receptor in mice  
 AU Cerwenka, Adelheid; Bakker, Alexander B. H.; McClanahan, Terri; Wagner, Janet; Wu, Jun; Phillips, Joseph H.; Lanier, Lewis L.  
 CS Department of Microbiology and Immunology and The Cancer Research Institute, University of California, San Francisco, San Francisco, CA, 94143, USA  
 SO Immunity (2000), 12(6), 721-727  
 CODEN: IUNIEH; ISSN: 1074-7613  
 PB Cell Press  
 DT Journal  
 LA English  
 AB Here we describe a family of GPI-anchored cell surface proteins that function as ligands for the mouse activating **NKG2D** receptor. These mols. are encoded by the retinoic acid early inducible (RAE-1) and H60 minor histocompatibility antigen genes on mouse chromosome 10 and show weak homol. with MHC class I. Expression of the **NKG2D** ligands is low or absent on normal, adult tissues; however, they are

constitutively expressed on some tumors and upregulated by retinoic acid. Ectopic expression of RAE-1 and H60 confers target susceptibility to NK cell attack. These studies identify a family of ligands for the activating **NKG2D** receptor on NK and T cells, which may play an important role in innate and adaptive immunity.

RE.CNT 32

RE

- (1) Aldrich, C; Cell 1994, V79, P649 HCAPLUS
  - (2) Bahram, S; Res Immunol 1996, V147, P328 HCAPLUS
  - (3) Bakker, A; Hum Immunol 2000, V61, P18 HCAPLUS
  - (4) Bauer, S; Science 1999, V285, P727 HCAPLUS
  - (5) Berg, S; Int Immunol 1998, V10, P379 HCAPLUS
- ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 15 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 2000:114379 BIOSIS

DN PREV200000114379

TI Paired inhibitory and triggering NK cell receptors for HLA class I molecules.

AU Lopez-Botet, Miguel (1); Bellon, Teresa; Llano, Manuel; Navarro, Francisco; Garcia, Pilar; de Miguel, Maria

CS (1) Servicio de Inmunologia, Hospital Universitario de la Princesa, Diego de Leon 62, 28006, Madrid Spain

SO Human Immunology, (Jan., 2000) Vol. 61, No. 1, pp. 7-17.  
ISSN: 0198-8859.

DT Article

LA English

SL English

AB Human natural killer (NK) cells specifically interact with major histocompatibility complex (MHC) class I molecules employing different receptor systems, shared with subsets of alphabeta and gammadelta T lymphocytes. Killer cell immunoglobulin-like receptors (KIRs) recognize groups of human leukocyte antigen (HLA) class Ia proteins displaying common structural features at the alpha-1 domain; among them, KIR2DL4 has been proposed to specifically interact with the class Ib molecule HLA-G1. Members of a related family of immunoglobulin (Ig)-like receptors (ILT2 or LIR-1 and ILT4 or LIR-2), expressed by other leukocyte lineages, interact with a broad spectrum of class Ia molecules and HLA-G1. On the other hand, CD94/NKG2-A(-C) and **NKG2D** lectin-like receptors, respectively, recognize the class Ib molecules HLA-E and MICA. A recurrent finding within the different receptor families is the existence of pairs of homologous molecules that often share the same **ligands** but display divergent functions. Inhibitory receptors tend to exhibit an affinity for HLA molecules higher than their activating counterparts. Recruitment of SH2 domain-bearing tyrosine phosphatases (SHP) by cytoplasmic phosphorylated immunoreceptor tyrosine-based inhibition motifs (ITIMs) is a crucial event for the inhibitory signalling pathway. By contrast, triggering receptors assemble with homodimers of immune tyrosine-based activation motif (ITAM)-bearing adaptor molecules (i.e., DAP12, CD3 zeta) that engage tyrosine kinases (ZAP70 and syk). Human Immunology 61, 7-17 (2000).

L22 ANSWER 16 OF 18 HCAPLUS COPYRIGHT 2001 ACS

AN 1999:765800 HCAPLUS

DN 132:62755

TI Natural killer cells: stress out, turn on, tune in

AU Diefenbach, Andreas; Raulet, David H.

CS Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California at Berkeley, Berkeley, CA, 94720-3200, USA

SO Curr. Biol. (1999), 9(22), R851-R853

CODEN: CUBLE2; ISSN: 0960-9822

PB Current Biology Publications

DT Journal; General Review



LA English  
 AB A review with 14 refs. Natural killer cells attack tumor cells, infected cells and some normal cells, but the basis of their specificity is not completely understood. Recent studies indicate that epithelial tumor cells upregulate a stress-induced MHC class-I-like protein termed MICA, triggering NK cells via a recently described receptor called **NKG2D**

RE.CNT 14

RE

- (1) Bahram, S; Proc Natl Acad Sci USA 1994, V91, P6259 HCAPLUS
- (2) Bauer, S; Science 1999, V285, P727 HCAPLUS
- (3) Correa, I; Eur J Immunol 1994, V24, P1323 HCAPLUS
- (4) Groh, V; Proc Natl Acad Sci USA 1996, V93, P12445 HCAPLUS
- (5) Groh, V; Proc Natl Acad Sci USA 1999, V96, P6879 HCAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L22 ANSWER 17 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1998:433707 BIOSIS

DN PREV199800433707

TI The genomic organization of NKG2C, E, F, and D receptor genes in the human natural killer gene complex.

AU Glienke, Jens; Sobanov, Yuri; Brostjan, Christine; Steffens, Christian; Nguyen, Catherine; Lehrach, Hans; Hofer, Erhard (1); Francis, Fiona

CS (1) Dep. Vascular Biol. Thrombosis Res., Univ. Vienna, Brunnerstr. 59, A-1235 Vienna Austria

SO Immunogenetics, (Aug., 1998) Vol. 48, No. 3, pp. 163-173.  
 ISSN: 0093-7711.

DT Article

LA English

AB Interactions of natural killer cell receptors with their cognate **ligands** play a major role in regulating NK cell function. The NKG2 gene family encodes several highly similar proteins, which are known to form heterodimers with the CD94 receptor. These dimers play a role in the inhibition as well as the activation of NK cells. We have analyzed the gene structures of the NKG2C, D, E, and F genes, and determined their genomic organization. Restriction mapping and sequencing revealed the four genes to be closely linked to one another, and of the same transcriptional orientation. An exon duplication within the NKG2C and E genes was identified, although the duplicated version of this exon has not yet been found in mRNA sequences. The NKG2C, E, and F genes, despite being highly similar, are variable at their 3' ends. We show that NKG2C consists of six exons, whereas NKG2E has seven, and the splice acceptor site for the seventh exon occurs in an Alu repeat. NKG2F consists of only four exons and part of exon IV is in some cases spliced to the 5' end of the **NKG2D** transcript. **NKG2D** has only a low similarity to the other NKG2 genes.

L22 ANSWER 18 OF 18 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1998:78276 BIOSIS

DN PREV199800078276

TI Cloning of a mouse homolog of CD94 extends the family of C-type lectins on murine natural killer cells.

AU Vance, Russell E.; Tanamachi, Dawn M.; Hanke, Thomas; Raulet, David H. (1)

CS (1) Dep. Mol. Cell Biol., Cancer Res. Lab., 485 LSA, Univ. Calif. at Berkeley, Berkeley, CA 94720 USA

SO European Journal of Immunology, (Dec., 1997) Vol. 27, No. 12, pp. 3236-3241.  
 ISSN: 0014-2980.

DT Article

LA English

AB Two families of major histocompatibility complex (MHC) class I-specific receptors are found on natural killer (NK) cells: immunoglobulin-like receptors and C-type lectin receptors. In mice, the latter category is

represented by the Ly49 family of receptors, whereas in humans, NK cells express the distantly related CD94, which forms MHC class I-specific heterodimers with NKG2 family members. Humans also express the MHC class I-specific p50/p58/p70 family of immunoglobulin-like receptors, but these have not been identified in mice. Hence, there is no known instance of an MHC class I-specific receptor that is expressed by both human and murine NK cells. Here we report the cloning of CD94 from the CB.17 and C57BL/6 strains of mice. Mouse CD94 is 54% identical and 66% similar to human CD94, and is also a member of the C-type lectin superfamily. Mouse CD94 is expressed efficiently on the cell surface of cells transiently transfected with the corresponding cDNA, but surface CD94 was unable to mediate detectable binding to MHC class I-expressing ConA blasts. Notably, mouse CD94, like human CD94, has a very short cytoplasmic tail, suggesting the existence of partner chains that may play a role in **ligand** binding and signaling. Like many other C-type lectins expressed by NK cells, mouse CD94 maps to the NK complex on distal chromosome 6, syntenic to human CD94. We also demonstrate that mouse CD94 is highly expressed specifically by mouse NK cells, raising the possibility that mice, like humans, express multiple families of MHC class I-specific receptors on their NK cells. Murine homologs of human NKG2 family members have not yet been identified, but we report here the existence of a murine **NKG2D**-like sequence that also maps to the murine NK complex near CD94 and Ly49 family members.

=> fil wpids

FILE 'WPIDS' ENTERED AT 13:55:42 ON 15 NOV 2001  
COPYRIGHT (C) 2001 DERWENT INFORMATION LTD

FILE LAST UPDATED: 13 NOV 2001 <20011113/UP>  
MOST RECENT DERWENT UPDATE 200166 <200166/DW>  
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> SDI'S MAY BE RUN ON EVERY UPDATE OR MONTHLY AS OF JUNE 2001.  
(EVERY UPDATE IS THE DEFAULT). FOR PRICING INFORMATION  
SEE HELP COST <<<

>>> FOR UP-TO-DATE INFORMATION ABOUT THE DERWENT CHEMISTRY  
RESOURCE, PLEASE VISIT  
<http://www.derwent.com/chemistryresource/index.html> <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES,  
SEE <http://www.derwent.com/dwpi/updates/dwpicov/index.html> <<<

=> d que 115

L15 1 SEA FILE=WPIDS ABB=ON NKG2D OR NKG2 D

=> d bib ab tech 115

L15 ANSWER 1 OF 1 WPIDS COPYRIGHT 2001 DERWENT INFORMATION LTD  
AN 1992-365992 [44] WPIDS  
DNC C1992-162471  
TI Isolated DNA or CDNA sequence encoding extracellular part of  
trans-membrane protein - used as immunosuppressants in organ transplants,  
and for treating auto-immune diseases, cancer and viral infections, also  
useful in diagnosis.  
DC B04 D16  
IN BACH, F H; HOFER, E; HOUCHINS, J P; MCSHERRY, C M; YABE, T S; YABE, T  
PA (SANO) SANDOZ LTD; (MINU) UNIV MINNESOTA; (NOVS) NOVARTIS AG  
CYC 17  
PI WO 9217198 A1 19921015 (199244)\* EN 62p  
RW: AT BE CH DE DK ES FR GB GR IT LU MC NL SE  
W: JP US  
EP 585257 A1 19940309 (199410) EN  
R: AT BE CH DE DK ES FR GB GR IT LI LU NL SE  
JP 06506358 W 19940721 (199433) 15p  
EP 585257 A4 19950222 (199611)  
US 6262244 B1 20010717 (200148)  
ADT WO 9217198 A1 WO 1992-US2469 19920327; EP 585257 A1 EP 1992-909331  
19920327, WO 1992-US2469 19920327; JP 06506358 W JP 1992-508930 19920327,  
WO 1992-US2469 19920327; EP 585257 A4 EP 1992-909331 ; US 6262244  
B1 CIP of US 1991-676663 19910328, Cont of WO 1992-US2469 19920327, Cont  
of US 1993-122514 19930924, US 1995-543246 19951013  
FDT EP 585257 A1 Based on WO 9217198; JP 06506358 W Based on WO 9217198  
PRAI US 1991-676663 19910328; US 1993-122514 19930924; US 1995-543246  
19951013  
AB WO 9217198 A UPAB: 19931116  
In an isolated DNA or cDNA encoding the extracellular part of a  
transmembrane protein designated (a) NKG2-A, (b) NKG2-B, (c) NKG2-C and  
(d) **NKG2-D** (f = fragment) translated in natural killer  
cells or T-cells, the DNA is selected from portions of DNA sequences in  
the specification.  
Also new are (a) isolated DNA or cDNA encoding a complete  
transmembrane protein designated NKG2-A NKG2-B, NKG2-C and **NKG2-D**  
(in specification); (b) isolated extracellular part, or the  
complete transmembrane protein (in specification), or sequence variants;

(c) poly- or monoclonal antibody recognising at least one epitope of (b);  
(d) bifunctional antibody recognising at least one epitope of (b) and a  
cancer- or virus-specific antigen; and (e) a chimeric protein mol.  
comprising (b) and a cytotoxic protein.

USE/ADVANTAGE - The DNA or chimeric protein may be used to treat a  
cancer or virus infection. Antibodies can activate natural killer cells  
and T-cells. The extracellular domains of the proteins are useful  
diagnostic tools for detecting target ligands, e.g. carbohydrate groups  
present on some cancer and virus-infected cells, and the complete protein  
may be used to study the mechanism of natural killer cell regulation.  
Dwg.0/1

Harris 09/871,491

=> fil medline

FILE 'MEDLINE' ENTERED AT 14:14:40 ON 15 NOV 2001

FILE LAST UPDATED: 14 NOV 2001 (20011114/UP). FILE COVERS 1958 TO DATE.

On April 22, 2001, MEDLINE was reloaded. See HELP RLOAD for details.

MEDLINE now contains IN-PROCESS records. See HELP CONTENT for details.

MEDLINE is now updated 4 times per week. A new current-awareness alert frequency (EVERYUPDATE) is available. See HELP UPDATE for more information.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2001 vocabulary. Enter HELP THESAURUS for details.

The OLDMEDLINE file segment now contains data from 1958 through 1965. Enter HELP CONTENT for details.

Left, right, and simultaneous left and right truncation are available in the Basic Index. See HELP SFIELDS for details.

THIS FILE CONTAINS CAS REGISTRY NUMBERS FOR EASY AND ACCURATE SUBSTANCE IDENTIFICATION.

=> d his

(FILE 'HOME' ENTERED AT 14:13:18 ON 15 NOV 2001)

FILE 'MEDLINE' ENTERED AT 14:13:24 ON 15 NOV 2001

L1 47 S NKG2D OR NKG2 D  
L2 23939 S LIGANDS/CT  
L3 101218 S RECEPTORS, IMMUNOLOGIC+NT/CT  
L4 11 S L1 AND L2  
L5 11 S L4 AND L3  
L6 546911 S TUMOR#  
L7 13 S L1 AND L6  
L8 7 S L7 NOT L5

FILE 'MEDLINE' ENTERED AT 14:14:40 ON 15 NOV 2001

=> d .med 15 1-11;d .med 18 1-7

L5 ANSWER 1 OF 11 MEDLINE  
AN 2001510266 MEDLINE  
DN 21441910 PubMed ID: 11557981  
TI Rael and H60 ligands of the NKG2D receptor stimulate tumour immunity.  
AU Diefenbach A; Jensen E R; Jamieson A M; Raulet D H  
CS Department of Molecular and Cell Biology and Cancer Research Laboratory, University of California, Berkeley 94720, USA.  
SO NATURE, (2001 Sep 13) 413 (6852) 165-71.  
Journal code: NSC; 0410462. ISSN: 0028-0836.  
CY England: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200110  
ED Entered STN: 20010917  
Last Updated on STN: 20011022  
Entered Medline: 20011018  
AB Natural killer (NK) cells attack many tumour cell lines, and are thought to have a critical role in anti-tumour immunity; however, the interaction

between NK cells and tumour targets is poorly understood. The stimulatory lectin-like **NKG2D** receptor is expressed by NK cells, activated CD8+ T cells and by activated macrophages in mice. Several distinct cell-surface ligands that are related to class I major histocompatibility complex molecules have been identified, some of which are expressed at high levels by tumour cells but not by normal cells in adults. However, no direct evidence links the expression of these 'induced self' ligands with tumour cell rejection. Here we demonstrate that ectopic expression of the murine **NKG2D** ligands Raelbeta or H60 in several tumour cell lines results in potent rejection of the tumour cells by syngeneic mice. Rejection is mediated by NK cells and/or CD8+ T cells. The ligand-expressing tumour cells induce potent priming of cytotoxic T cells and sensitization of NK cells in vivo. Mice that are exposed to live or irradiated tumour cells expressing Rael or H60 are specifically immune to subsequent challenge with tumour cells that lack **NKG2D** ligands, suggesting application of the ligands in the design of tumour vaccines.

CT Check Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

\*CD8-Positive T-Lymphocytes: IM, immunology

Cytotoxicity, Immunologic

Immunity

\*Killer Cells, Natural: IM, immunology

#### Ligands

\*Membrane Proteins: ME, metabolism

Mice

Mice, Inbred C57BL

\*Minor Histocompatibility Antigens: ME, metabolism

\*Neoplasms: IM, immunology

\*Receptors, Immunologic: ME, metabolism

Recombinant Proteins

T-Lymphocytes, Cytotoxic: IM, immunology

Tumor Cells, Cultured

L5 ANSWER 2 OF 11 MEDLINE

AN 2001445970 MEDLINE

DN 21383614 PubMed ID: 11491531

TI Interactions of human **NKG2D** with its ligands MICA, MICB, and homologs of the mouse RAE-1 protein family.

AU Steinle A; Li P; Morris D L; Groh V; Lanier L L; Strong R K; Spies T

CS Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle WA 98109, USA.

NC AI30581 (NIAID)

AI42200 (NIAID)

SO IMMUNOGENETICS, (2001 May-Jun) 53 (4) 279-87.

Journal code: GI4; 0420404. ISSN: 0093-7711.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200108

ED Entered STN: 20010813

Last Updated on STN: 20010827

Entered Medline: 20010823

AB **NKG2D** is an activating receptor that is expressed on most natural killer (NK) cells, CD8 alphabeta T cells, and gammadelta T cells. Among its ligands is the distant major histocompatibility complex class I homolog MICA, which has no function in antigen presentation but is induced by cellular stress. To extend previous functional evidence, the **NKG2D**-MICA interaction was studied in isolation. **NKG2D** homodimers formed stable complexes with monomeric MICA in solution, demonstrating that no other components were required to facilitate this interaction. MICA glycosylation was not essential but enhanced complex formation. Soluble **NKG2D** also bound to cell surface MICB, which has structural and functional properties similar to those of MICA.

Moreover, **NKG2D** stably interacted with surface molecules encoded by three newly identified cDNA sequences (N2DL-1, -2, and -3), which are identical to the human ULBP proteins and may represent homologs of the mouse retinoic acid-early inducible family of **NKG2D** ligands. Because of the substantial sequence divergence among these molecules, these results indicated promiscuous modes of receptor binding. Comparison of allelic variants of MICA revealed large differences in **NKG2D** binding that were associated with a single amino acid substitution at position 129 in the alpha2 domain. Varying affinities of MICA alleles for **NKG2D** may affect thresholds of NK-cell triggering and T-cell modulation.

CT Check Tags: Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.  
Amino Acid Sequence

Antigens, Surface: ME, metabolism

Dimerization

Histocompatibility Antigens Class I: CH, chemistry

\*Histocompatibility Antigens Class I: ME, metabolism

\*Killer Cells, Natural: IM, immunology

#### Ligands

\*Membrane Proteins: ME, metabolism

Models, Molecular

Molecular Sequence Data

Protein Binding

\*Receptors, Immunologic: ME, metabolism

Sequence Homology, Amino Acid

Solubility

L5 ANSWER 3 OF 11 MEDLINE

AN 2001373026 MEDLINE

DN 21323007 PubMed ID: 11429322

TI MICA and MICB genes: can the enigma of their polymorphism be resolved?.

AU Stephens H A

CS Institute of Urology and Nephrology, University College London, The Middlesex Hospital, 48 Riding House Street, London, UK, W1W 7EY..  
h.stephens@ucl.ac.uk

SO Trends Immunol, (2001 Jul) 22 (7) 378-85. Ref: 72

Journal code: DZX; 100966032. ISSN: 1471-4906.

CY England: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

General Review; (REVIEW)

(REVIEW, TUTORIAL)

LA English

FS Priority Journals

EM 200108

ED Entered STN: 20010903

Last Updated on STN: 20010903

Entered Medline: 20010830

AB The human MHC class I chain-related genes (MICA and MICB) are located within the HLA class I region of chromosome 6. Their organization, expression and products differ considerably from classical HLA class I genes. MIC proteins are considered to be markers of "stress" in the epithelia, and act as ligands for cells expressing a common activatory natural killer-cell receptor (**NKG2D**). Molecular models are now available for the MICA protein, both bound and complexed with **NKG2D**. MICA molecules appear to be highly flexible and polymorphic, although the functional relevance and implications of their polymorphism have yet to be fully discerned.

CT Check Tags: Animal; Human; Support, Non-U.S. Gov't

Alleles

Chromosome Mapping

Epithelial Cells

Evolution, Molecular

Gene Expression

Genetics, Population  
 \*Histocompatibility Antigens Class I: GE, genetics  
 Histocompatibility Antigens Class I: IM, immunology  
 Killer Cells, Natural: IM, immunology

**Ligands**

\*Polymorphism (Genetics)  
**Receptors, Immunologic: IM, immunology**

L5 ANSWER 4 OF 11 MEDLINE  
 AN 2001309272 MEDLINE  
 DN 21223631 PubMed ID: 11323699  
 TI Complex structure of the activating immunoreceptor **NKG2D** and its MHC class I-like ligand MICA.  
 CM Comment in: Nat Immunol. 2001 May;2(5):379-80  
 AU Li P; Morris D L; Willcox B E; Steinle A; Spies T; Strong R K  
 CS Division of Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA 98109 USA.  
 NC AI30581 (NIAID)  
 AI42200 (NIAID)  
 CA18221 (NCI)  
 SO Nat Immunol, (2001 May) 2 (5) 443-51.  
 Journal code: DOG; 100941354. ISSN: 1529-2908.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 200105  
 ED Entered STN: 20010604  
 Last Updated on STN: 20010604  
 Entered Medline: 20010531  
 AB The major histocompatibility complex (MHC) class I homolog, MICA, is a stress-inducible ligand for **NKG2D**, a C-type lectin-like activating immunoreceptor. The crystal structure of this ligand-receptor complex that we report here reveals an **NKG2D** homodimer bound to a MICA monomer in an interaction that is analogous to that seen in T cell receptor-MHC class I protein complexes. Similar surfaces on each **NKG2D** monomer interact with different surfaces on either the alpha1 or alpha2 domains of MICA. The binding interactions are large in area and highly complementary. The central section of the alpha2-domain helix, disordered in the structure of MICA alone, is ordered in the complex and forms part of the **NKG2D** interface. The extensive flexibility of the interdomain linker of MICA is shown by its altered conformation when crystallized alone or in complex with **NKG2D**.  
 CT Check Tags: Human; Support, U.S. Gov't, P.H.S.  
 Amino Acid Sequence  
 \*Histocompatibility Antigens Class I: CH, chemistry  
 \*Killer Cells, Natural: IM, immunology  
 Lectins: CH, chemistry  
**Ligands**  
 Models, Molecular  
 Molecular Sequence Data  
 Protein Binding  
 Protein Conformation  
**\*Receptors, Immunologic: CH, chemistry**  
 Sequence Homology, Amino Acid  
 Surface Plasmon Resonance  
 Surface Properties  
 L5 ANSWER 5 OF 11 MEDLINE  
 AN 2001216034 MEDLINE  
 DN 21205390 PubMed ID: 11248803  
 TI Ligands for the murine **NKG2D** receptor: expression by tumor cells and activation of NK cells and macrophages.



CM Comment in: Nat Immunol. 2000 Aug;1(2):95-7  
 AU Diefenbach A; Jamieson A M; Liu S D; Shastri N; Raulet D H  
 CS Department of Molecular and Cell Biology and Cancer Research Laboratory,  
 University of California, Berkeley, USA.  
 SO Nat Immunol, (2000 Aug) 1 (2) 119-26.  
 Journal code: DOG; 100941354. ISSN: 1529-2908.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 200105  
 ED Entered STN: 20010521  
 Last Updated on STN: 20010521  
 Entered Medline: 20010517  
 AB Natural killer (NK) cells attack tumor and infected cells, but the  
 receptors and ligands that stimulate them are poorly understood. Here we  
 report the expression cloning of two murine ligands for the lectin-like  
 receptor **NKG2D**. The two ligands, H-60 and Rael beta, are distant  
 relatives of major histocompatibility complex class I molecules.  
**NKG2D** ligands are not expressed by most normal cells but are  
 up-regulated on numerous tumor cells. We show that mouse **NKG2D**  
 is expressed by NK cells, activated CD8+ T cells and activated  
 macrophages. Expression of either **NKG2D** ligand by target cells  
 triggers NK cell cytotoxicity and interferon-gamma secretion by NK cells,  
 as well as nitric oxide release and tumor necrosis factor alpha  
 transcription by macrophages. Thus, through their interaction with  
**NKG2D**, H-60 and Rael beta are newly identified potent stimulators  
 of innate immunity.  
 CT Check Tags: Animal; Human; Support, Non-U.S. Gov't; Support, U.S. Gov't,  
 P.H.S.  
 CHO Cells  
 COS Cells  
 Cercopithecus aethiops  
 Cloning, Molecular  
 Hamsters  
 \*Killer Cells, Natural: IM, immunology  
 Killer Cells, Natural: ME, metabolism  
**Ligands**  
 Lymphocyte Transformation  
 Macrophage Activation  
 \*Macrophages, Peritoneal: IM, immunology  
 Macrophages, Peritoneal: ME, metabolism  
 Membrane Proteins: GE, genetics  
 \*Membrane Proteins: IM, immunology  
 Mice  
 Mice, Inbred BALB C  
 Mice, Inbred C57BL  
 Minor Histocompatibility Antigens: GE, genetics  
 \*Minor Histocompatibility Antigens: IM, immunology  
**Receptors, Immunologic: GE, genetics**  
**\*Receptors, Immunologic: ME, metabolism**  
 Tumor Cells, Cultured  
 L5 ANSWER 6 OF 11 MEDLINE  
 AN 2001210233 MEDLINE  
 DN 21195294 PubMed ID: 11298332  
 TI Role of **NKG2D** in tumor cell lysis mediated by human NK cells:  
 cooperation with natural cytotoxicity receptors and capability of  
 recognizing tumors of nonepithelial origin.  
 AU Pende D; Cantoni C; Rivera P; Vitale M; Castriconi R; Marcenaro S; Nanni  
 M; Biassoni R; Bottino C; Moretta A; Moretta L  
 CS Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy.  
 SO EUROPEAN JOURNAL OF IMMUNOLOGY, (2001 Apr) 31 (4) 1076-86.

Journal code: EN5; 1273201. ISSN: 0014-2980.

CY Germany: Germany, Federal Republic of

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200105

ED Entered STN: 20010517

Last Updated on STN: 20010517

Entered Medline: 20010510

AB **NKG2D** is a recently described activating receptor expressed by both NK cells and CTL. In this study we investigated the role of **NKG2D** in the natural cytotoxicity mediated by NK cell clones. The role of **NKG2D** varied depending on the type of target cells analyzed. Lysis of various tumors appeared to be exclusively natural cytotoxicity receptors (NCR) dependent. In contrast, killing of another group of target cells, including not only the epithelial cell lines HELA and IGROV-1, but also the FO-1 melanoma, the JA3 leukemia, the Daudi Burkitt lymphoma and even normal PHA-induced lymphoblasts, involved both NCR and **NKG2D**. Notably, NK cell clones expressing low surface densities of NCR (NCR(dull)) could lyse these tumors in an exclusively **NKG2D**-dependent fashion. Remarkably, not all of these targets expressed MICA/B, thus implying the existence of additional ligands recognized by **NKG2D**, possibly represented by GPI-linked molecules. Finally, we show that the engagement of different HLA class I-specific inhibitory receptors by either specific antibodies or the appropriate HLA class I ligand led to inhibition of **NKG2D**-mediated NK cell triggering.

CT Check Tags: Animal; Human; Support, Non-U.S. Gov't

Antibodies, Monoclonal: IM, immunology

Antibodies, Monoclonal: PD, pharmacology

Cells, Cultured

Clone Cells: DE, drug effects

Clone Cells: IM, immunology

\*Cytotoxicity, Immunologic

Cytotoxicity, Immunologic: DE, drug effects

Down-Regulation (Physiology)

Epithelial Cells: IM, immunology

Epithelial Cells: PA, pathology

Flow Cytometry

Histocompatibility Antigens Class I: IM, immunology

Killer Cells, Natural: DE, drug effects

\*Killer Cells, Natural: IM, immunology

Killer Cells, Natural: ME, metabolism

#### Ligands

Mice

\*Neoplasms: IM, immunology

\*Neoplasms: PA, pathology

Phytohemagglutinins: IM, immunology

Phytohemagglutinins: PD, pharmacology

RNA, Messenger: AN, analysis

RNA, Messenger: GE, genetics

Receptors, Immunologic: GE, genetics

\*Receptors, Immunologic: IM, immunology

Transfection

Tumor Cells, Cultured

L5 ANSWER 7 OF 11 MEDLINE

AN 2001205813 MEDLINE

DN 21137928 PubMed ID: 11239445

TI ULBPs, novel MHC class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the **NKG2D** receptor.

AU Cosman D; Mullberg J; Sutherland C L; Chin W; Armitage R; Fanslow W; Kubin M; Chalupny N J

CS Department of Molecular Biology, Immunex Corporation, 51 University  
Street, Seattle, WA 98101, . USA.cosman@immunex.com

SO IMMUNITY, (2001 Feb) 14 (2) 123-33.  
Journal code: CCF; 9432918. ISSN: 1074-7613.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

OS GENBANK-AF304377; GENBANK-AF304378; GENBANK-AF304379

EM 200104

ED Entered STN: 20010417  
Last Updated on STN: 20010417  
Entered Medline: 20010412

AB The human cytomegalovirus glycoprotein, UL16, binds to two members of a  
novel family of molecules, the ULBPs, and to the MHC class I homolog,  
MICB. The ULBPs are GPI-linked glycoproteins belonging to the extended MHC  
class I family but are only distantly related to MICB. The ULBP and MICB  
molecules are ligands for the activating receptor, **NKG2D**/DAPI10,  
and this interaction is blocked by a soluble form of UL16. The ULBPs  
stimulate cytokine and chemokine production from NK cells, and expression  
of ULBPs in NK cell-resistant target cells confers susceptibility to NK  
cell cytotoxicity. Masking of NK cell recognition of ULBP or MIC antigens  
by UL16 provides a potential mechanism by which human cytomegalovirus-  
infected cells might evade attack by the immune system.

CT Check Tags: Human  
Amino Acid Sequence  
Base Sequence  
Carrier Proteins: GE, genetics  
\*Carrier Proteins: IM, immunology  
\*Carrier Proteins: ME, metabolism  
Cell Line  
\*Cytomegalovirus: IM, immunology  
\*Cytomegalovirus: ME, metabolism  
Cytomegalovirus: PY, pathogenicity  
Cytotoxicity, Immunologic  
DNA Primers: GE, genetics  
Glycoproteins: IM, immunology  
Glycoproteins: ME, metabolism  
Histocompatibility Antigens Class I: GE, genetics  
\*Histocompatibility Antigens Class I: IM, immunology  
\*Histocompatibility Antigens Class I: ME, metabolism  
\*Killer Cells, Natural: IM, immunology

**Ligands**  
Molecular Sequence Data  
\*Receptors, Immunologic: ME, metabolism  
Sequence Homology, Amino Acid  
\*Viral Proteins: IM, immunology  
\*Viral Proteins: ME, metabolism

L5 ANSWER 8 OF 11 MEDLINE

AN 2000350669 MEDLINE

DN 20350669 PubMed ID: 10894171

TI Retinoic acid early inducible genes define a ligand family for the  
activating **NKG2D** receptor in mice.

AU Cerwenka A; Bakker A B; McClanahan T; Wagner J; Wu J; Phillips J H; Lanier  
L L

CS Department of Microbiology and Immunology and The Cancer Research  
Institute, University of California, San Francisco 94143, USA.

SO IMMUNITY, (2000 Jun) 12 (6) 721-7.  
Journal code: CCF; 9432918. ISSN: 1074-7613.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals  
 OS GENBANK-AF257520  
 EM 200007  
 ED Entered STN: 20000811  
 Last Updated on STN: 20000811  
 Entered Medline: 20000731

AB Here we describe a family of GPI-anchored cell surface proteins that function as ligands for the mouse activating **NKG2D** receptor. These molecules are encoded by the retinoic acid early inducible (RAE-1) and H60 minor histocompatibility antigen genes on mouse chromosome 10 and show weak homology with MHC class I. Expression of the **NKG2D** ligands is low or absent on normal, adult tissues; however, they are constitutively expressed on some tumors and upregulated by retinoic acid. Ectopic expression of RAE-1 and H60 confers target susceptibility to NK cell attack. These studies identify a family of ligands for the activating **NKG2D** receptor on NK and T cells, which may play an important role in innate and adaptive immunity.

CT Check Tags: Animal; Human; Support, Non-U.S. Gov't  
 Amino Acid Sequence  
 Carcinoma, Lewis Lung  
 Cloning, Molecular: MT, methods  
 Cytotoxicity, Immunologic: DE, drug effects  
 Gene Expression Regulation: DE, drug effects  
 \*Gene Expression Regulation: IM, immunology  
 Glycosylphosphatidylinositols: ME, metabolism  
 IgG: GE, genetics  
 Immunoglobulins, Fc: GE, genetics  
 Killer Cells, Natural: IM, immunology  
 \*Killer Cells, Natural: ME, metabolism

**Ligands**  
 Membrane Proteins: BI, biosynthesis  
 \*Membrane Proteins: GE, genetics  
 Membrane Proteins: ME, metabolism  
 Membrane Proteins: PH, physiology  
 Mice  
 Mice, Inbred C57BL  
 Minor Histocompatibility Antigens: BI, biosynthesis  
 Minor Histocompatibility Antigens: GE, genetics  
 Minor Histocompatibility Antigens: ME, metabolism  
 Minor Histocompatibility Antigens: PH, physiology  
 Molecular Sequence Data  
 \*Multigene Family: IM, immunology

**Receptors, Immunologic: GE, genetics**  
**\*Receptors, Immunologic: ME, metabolism**  
 Recombinant Fusion Proteins: GE, genetics  
 Recombinant Fusion Proteins: ME, metabolism  
 \*Tretinoin: PD, pharmacology  
 Tumor Cells, Cultured

L5 ANSWER 9 OF 11 MEDLINE  
 AN 2000122083 MEDLINE  
 DN 20122083 PubMed ID: 10658973  
 TI Paired inhibitory and triggering NK cell receptors for HLA class I molecules.  
 AU Lopez-Botet M; Bellon T; Llano M; Navarro F; Garcia P; de Miguel M  
 CS Servicio de Immunologia, Hospital Universitario de la Princesa, Madrid, Spain.. mlbotet@hup.es  
 SO HUMAN IMMUNOLOGY, (2000 Jan) 61 (1) 7-17. Ref: 108  
 Journal code: G9W; 8010936. ISSN: 0198-8859.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)

LA English  
 FS Priority Journals  
 EM 200003  
 ED Entered STN: 20000314  
 Last Updated on STN: 20000314  
 Entered Medline: 20000302

AB Human natural killer (NK) cells specifically interact with major histocompatibility complex (MHC) class I molecules employing different receptor systems, shared with subsets of alphabeta and gammadelta T lymphocytes. Killer cell immunoglobulin-like receptors (KIRs) recognize groups of human leukocyte antigen (HLA) class Ia proteins displaying common structural features at the alpha-1 domain; among them, KIR2DL4 has been proposed to specifically interact with the class Ib molecule HLA-G1. Members of a related family of immunoglobulin (Ig)-like receptors (ILT2 or LIR-1 and ILT4 or LIR-2), expressed by other leukocyte lineages, interact with a broad spectrum of class Ia molecules and HLA-G1. On the other hand, CD94/NKG2-A(-C) and **NKG2D** lectin-like receptors, respectively, recognize the class Ib molecules HLA-E and MICA. A recurrent finding within the different receptor families is the existence of pairs of homologous molecules that often share the same ligands but display divergent functions. Inhibitory receptors tend to exhibit an affinity for HLA molecules higher than their activating counterparts. Recruitment of SH2 domain-bearing tyrosine phosphatases (SHP) by cytoplasmic phosphorylated immunoreceptor tyrosine-based inhibition motifs (ITIMs) is a crucial event for the inhibitory signalling pathway. By contrast, triggering receptors assemble with homodimers of immune tyrosine-based activation motif (ITAM)-bearing adaptor molecules (i.e., DAP12, CD3 xi) that engage tyrosine kinases (ZAP70 and syk).

CT Check Tags: Human  
 Histocompatibility Antigens Class I: IM, immunology  
 \*Histocompatibility Antigens Class I: ME, metabolism  
 \*Killer Cells, Natural: IM, immunology  
 Leukocytes: IM, immunology  
 Ligands  
 Receptors, Immunologic: CL, classification  
 Receptors, Immunologic: GE, genetics  
 Receptors, Immunologic: IM, immunology  
 \*Receptors, Immunologic: ME, metabolism

L5 ANSWER 10 OF 11 MEDLINE  
 AN 1999357865 MEDLINE  
 DN 99357865 PubMed ID: 10426994  
 TI An activating immunoreceptor complex formed by **NKG2D** and DAP10.  
 CM Comment in: Science. 1999 Jul 30;285(5428):645-6  
 AU Wu J; Song Y; Bakker A B; Bauer S; Spies T; Lanier L L; Phillips J H  
 CS DNAX Research Institute, 901 California Avenue, Palo Alto, CA 94304, USA.  
 NC AI30581 (NIAID)  
 SO SCIENCE, (1999 Jul 30) 285 (5428) 730-2.  
 Journal code: UJ7; 0404511. ISSN: 0036-8075.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 OS GENBANK-AF072844; GENBANK-AF072845; GENBANK-AF072846; GENBANK-AF122904;  
 GENBANK-AF122905  
 EM 199908  
 ED Entered STN: 19990827  
 Last Updated on STN: 19990827  
 Entered Medline: 19990816

AB Many immune receptors are composed of separate ligand-binding and signal-transducing subunits. In natural killer (NK) and T cells, DAP10 was identified as a cell surface adaptor protein in an activating receptor complex with **NKG2D**, a receptor for the stress-inducible and

tumor-associated major histocompatibility complex molecule MICA. Within the DAP10 cytoplasmic domain, an Src homology 2 (SH2) domain-binding site was capable of recruiting the p85 subunit of the phosphatidylinositol 3-kinase (PI 3-kinase), providing for **NKG2D**-dependent signal transduction. Thus, **NKG2D**-DAP10 receptor complexes may activate NK and T cell responses against MICA-bearing tumors.

CT Check Tags: Animal; Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

1-Phosphatidylinositol 3-Kinase: ME, metabolism

Amino Acid Sequence

Binding Sites

Cell Line

Cytotoxicity, Immunologic

\*Killer Cells, Natural: IM, immunology

Killer Cells, Natural: ME, metabolism

**Ligands**

\*Lymphocyte Transformation

Membrane Proteins: CH, chemistry

Membrane Proteins: GE, genetics

\*Membrane Proteins: ME, metabolism

Mice

Molecular Sequence Data

Neoplasms: IM, immunology

Phosphorylation

Phosphotyrosine: ME, metabolism

Receptors, Immunologic: CH, chemistry

Receptors, Immunologic: GE, genetics

\*Receptors, Immunologic: ME, metabolism

Signal Transduction

\*T-Lymphocytes: IM, immunology

T-Lymphocytes: ME, metabolism

Tumor Cells, Cultured

src Homology Domains

L5 ANSWER 11 OF 11 MEDLINE

AN 1999357864 MEDLINE

DN 99357864 PubMed ID: 10426993

TI Activation of NK cells and T cells by **NKG2D**, a receptor for stress-inducible MICA.

CM Comment in: Science. 1999 Jul 30;285(5428):645-6

AU Bauer S; Groh V; Wu J; Steinle A; Phillips J H; Lanier L L; Spies T

CS Fred Hutchinson Cancer Research Center, Clinical Research Division, 1100 Fairview Avenue North, Seattle, WA 98109, USA.

NC P01 CA18221 (NCI)

RO1 AI30581 (NIAID)

SO SCIENCE, (1999 Jul 30) 285 (5428) 727-9.

Journal code: UJ7; 0404511. ISSN: 0036-8075.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199908

ED Entered STN: 19990827

Last Updated on STN: 19990827

Entered Medline: 19990816

AB Stress-inducible MICA, a distant homolog of major histocompatibility complex (MHC) class I, functions as an antigen for gammadelta T cells and is frequently expressed in epithelial tumors. A receptor for MICA was detected on most gammadelta T cells, CD8+ alphabeta T cells, and natural killer (NK) cells and was identified as **NKG2D**. Effector cells from all these subsets could be stimulated by ligation of **NKG2D**. Engagement of **NKG2D** activated cytolytic responses of gammadelta T cells and NK cells against transfectants and epithelial tumor cells

expressing MICA. These results define an activating immunoreceptor-MHC ligand interaction that may promote antitumor NK and T cell responses.

CT Check Tags: Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S. Cytotoxicity, Immunologic

\*Histocompatibility Antigens Class I: IM, immunology  
Histocompatibility Antigens Class I: ME, metabolism  
Jurkat Cells

\*Killer Cells, Natural: IM, immunology

**Ligands**  
Lymphocyte Subsets: IM, immunology

\*Lymphocyte Transformation  
Membrane Proteins: ME, metabolism

Receptors, Antigen, T-Cell, gamma-delta: IM, immunology  
Receptors, Immunologic: CH, chemistry  
Receptors, Immunologic: GE, genetics  
\*Receptors, Immunologic: IM, immunology  
Receptors, Immunologic: ME, metabolism

Signal Transduction  
\*T-Lymphocytes: IM, immunology  
Transfection  
Tumor Cells, Cultured

L8 ANSWER 1 OF 7 MEDLINE  
AN 2001563617 IN-PROCESS  
DN 21521655 PubMed ID: 11567108  
TI Immunology: stress, nk receptors, and immune surveillance.  
AU Pardoll D M  
SO SCIENCE, (2001 Oct 19) 294 (5542) 534-6.  
Journal code: UJ7; 0404511. ISSN: 0036-8075.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS IN-PROCESS; NONINDEXED; Priority Journals  
ED Entered STN: 20011022  
Last Updated on STN: 20011022

AB It has long been suspected that natural killer (NK) cells are involved in immune surveillance. Now comes new work, described by Pardoll in his Perspective, showing that gd T cells expressing an NK cell receptor called **NKG2d** are important for detecting precancerous skin epithelial cells in mice (Girardi et al.). Engagement of **NKG2d** with its ligands Rae-1 or H60 expressed on mouse epidermal cells treated with carcinogens results in activation of gd T cells, which eliminate the precancerous epidermal cells before they become established as tumors.

L8 ANSWER 2 OF 7 MEDLINE  
AN 2001525841 IN-PROCESS  
DN 21457265 PubMed ID: 11562472  
TI Ectopic expression of retinoic acid early inducible-1 gene (RAE-1) permits natural killer cell-mediated rejection of a MHC class I-bearing tumor in vivo.  
AU Cerwenka A; Baron J L; Lanier L L  
CS Department of Microbiology and Immunology and the Cancer Research Institute, University of California, San Francisco, CA 94143-0414.  
SO PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (2001 Sep 25) 98 (20) 11521-6.  
Journal code: PV3; 7505876. ISSN: 0027-8424.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English

FS IN-PROCESS; NONINDEXED; Priority Journals  
 ED Entered STN: 20010927  
 Last Updated on STN: 20010927

AB In 1986, Karre and colleagues reported that natural killer (NK) cells rejected an MHC class I-deficient **tumor** cell line (RMA-S) but they did not reject the same cell line if it expressed MHC class I (RMA). Based on this observation, they proposed the concept that NK cells provide immune surveillance for "missing self," e.g., they eliminate cells that have lost class I MHC antigens. This seminal observation predicted the existence of inhibitory NK cell receptors for MHC class I. Here, we present evidence that NK cells are able to reject **tumors** expressing MHC class I if the **tumor** expresses a ligand for **NKG2D**. Mock-transfected RMA cells resulted in **tumor** formation. In contrast, when RMA cells were transfected with the retinoic acid early inducible gene-1 gamma or delta (RAE-1), ligands for the activating receptor **NKG2D**, the **tumors** were rejected. The **tumor** rejection was mediated by NK cells, and not by CD1-restricted NK1.1(+) T cells. No T cell-mediated immunological memory against the parental **tumor** was generated in the animals that had rejected the RAE-1 transfected **tumors**, which succumbed to rechallenge with the parental RMA **tumor**. Therefore, NK cells are able to reject a **tumor** expressing RAE-1 molecules, despite expression of self MHC class I on the **tumor**, demonstrating the potential for NK cells to participate in immunity against class I-bearing malignancies.

L8 ANSWER 3 OF 7 MEDLINE  
 AN 2001468526 IN-PROCESS  
 DN 21404051 PubMed ID: 11513139  
 TI The UL16-binding proteins, a novel family of MHC class I-related ligands for **NKG2D**, activate natural killer cell functions.  
 AU Sutherland C L; Chalupny N J; Cosman D  
 CS Department of Molecular Biology, Immunex Corporation, Seattle, Washington 98101, USA.. sutherlandc@immunex.com  
 SO IMMUNOLOGICAL REVIEWS, (2001 Jun) 181 185-92.  
 Journal code: GG4; 7702118. ISSN: 0105-2896.  
 CY Denmark  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS IN-PROCESS; NONINDEXED; Priority Journals  
 ED Entered STN: 20010830  
 Last Updated on STN: 20010830

AB The UL16-binding proteins (ULBPs) are a novel family of MHC class I-related molecules (MICs) that were identified based on their ability to bind to the human cytomegalovirus (HCMV) glycoprotein UL16. UL16 also binds to a member of another family of MHC class I-like molecules, MICB. The ULBPs and MICs are ligands for **NKG2D**/DAP10, an activating receptor expressed by natural killer (NK) cells and other immune effector cells, and this interaction can be blocked by UL16. Engagement of **NKG2D**/DAP10 by ULBPs or MICs expressed on a target cell can overcome an inhibitory signal generated by NK-cell recognition of MHC class I molecules and trigger NK cytotoxicity. ULBPs elicit their effects on NK cells by activating the janus kinase 2, signal transducer and activator of transcription 5, extracellular-signal-regulated kinase mitogen-activated protein kinase and Akt/protein kinase B signal transduction pathways. Although ULBPs alone activate multiple signaling pathways and induce modest cytokine production, ULBPs synergize strongly with interleukin-12 for production of interferon-gamma by NK cells. This finding is consistent with reports in T cells that **NKG2D**/DAP10 can act as a co-stimulatory receptor in a similar manner as CD28. The possible roles of ULBPs in mediating immune responses to viruses and **tumors** and the potential mechanisms by which UL16 may allow HCMV to evade immune detection are areas of active investigation.



L8 ANSWER 4 OF 7 MEDLINE  
 AN 2001434402 MEDLINE  
 DN 21139817 PubMed ID: 11244035  
 TI Activating receptors and coreceptors involved in human natural killer cell-mediated cytotoxicity.  
 AU Moretta A; Bottino C; Vitale M; Pende D; Cantoni C; Mingari M C; Biassoni R; Moretta L  
 CS Dipartimento di Medicina Sperimentale, Universita degli Studi di Genova, Italy.. alemoret@unige.it  
 SO ANNUAL REVIEW OF IMMUNOLOGY, (2001) 19 197-223. Ref: 127  
 Journal code: ALO; 8309206. ISSN: 0732-0582.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, ACADEMIC)  
 LA English  
 FS Priority Journals  
 EM 200108  
 ED Entered STN: 20010806  
 Last Updated on STN: 20010806  
 Entered Medline: 20010802  
 AB Natural killer cells can discriminate between normal cells and cells that do not express adequate amounts of major histocompatibility complex (MHC) class I molecules. The discovery, both in mouse and in human, of MHC-specific inhibitory receptors clarified the molecular basis of this important NK cell function. However, the triggering receptors responsible for positive NK cell stimulation remained elusive until recently. Some of these receptors have now been identified in humans, thus shedding some light on the molecular mechanisms involved in NK cell activation during the process of natural cytotoxicity. Three novel, NK-specific, triggering surface molecules (NKp46, NKp30, and NKp44) have been identified. They represent the first members of a novel emerging group of receptors collectively termed natural cytotoxicity receptors (NCR). Monoclonal antibodies (mAbs) to NCR block to differing extents the NK-mediated lysis of various **tumors**. Moreover, lysis of certain **tumors** can be virtually abrogated by the simultaneous masking of the three NCRs. There is a coordinated surface expression of the three NCRs, their surface density varying in different individuals and also in the NK cells isolated from a given individual. A direct correlation exists between the surface density of NCR and the ability of NK cells to kill various **tumors**. NKp46 is the only NCR involved in human NK-mediated killing of murine target cells. Accordingly, a homologue of NKp46 has been detected in mouse. Molecular cloning of NCR revealed novel members of the Ig superfamily displaying a low degree of similarity to each other and to known human molecules. NCRs are coupled to different signal transducing adaptor proteins, including CD3 zeta, Fc epsilon RI gamma, and KARAP/DAP12. Another triggering NK receptor is **NKG2D**. It appears to play either a complementary or a synergistic role with NCRs. Thus, the triggering of NK cells in the process of **tumor** cell lysis may often depend on the concerted action of NCR and **NKG2D**. In some instances, however, it may uniquely depend upon the activity of NCR or **NKG2D** only. Strict **NKG2D**-dependency can be appreciated using clones that, in spite of their NCR(dull) phenotype, efficiently lyse certain epithelial **tumors** or leukemic cell lines. Other triggering surface molecules including 2B4 and the novel NKp80 appear to function as coreceptors rather than as true receptors. Indeed, they can induce natural cytotoxicity only when co-engaged with a triggering receptor. While an altered expression or function of NCR or **NKG2D** is being explored as a possible cause of immunological disorders, 2B4 dysfunction has already been associated with a severe form of immunodeficiency. Indeed, in patients with the X-linked lymphoproliferative disease, the inability to control Epstein-Barr virus

infections may be consequent to a major dysfunction of 2B4 that exerts inhibitory instead of activating functions.

CT Check Tags: Animal; Human; Support, Non-U.S. Gov't  
 Antibodies, Monoclonal: IM, immunology  
 Antibodies, Monoclonal: PD, pharmacology  
 Carrier Proteins: IM, immunology  
 Cloning, Molecular  
 \*Cytotoxicity, Immunologic: IM, immunology  
 Epstein-Barr Virus Infections: IM, immunology  
 Histocompatibility Antigens Class I: IM, immunology  
 \*Killer Cells, Natural: IM, immunology  
 Lymphoproliferative Disorders: IM, immunology  
 Membrane Glycoproteins: CH, chemistry  
 Membrane Glycoproteins: IM, immunology  
 Mice  
 Multigene Family  
 Neoplasms: IM, immunology  
 Neoplasms, Experimental: IM, immunology  
 Receptors, Immunologic: AI, antagonists & inhibitors  
 Receptors, Immunologic: CH, chemistry  
 \*Receptors, Immunologic: IM, immunology  
 Signal Transduction

L8 ANSWER 5 OF 7 MEDLINE  
 AN 2001106858 MEDLINE  
 DN 20578924 PubMed ID: 11137207  
 TI Triggering receptors involved in natural killer cell-mediated cytotoxicity against choriocarcinoma cell lines.  
 AU Sivori S; Parolini S; Marcenaro E; Millo R; Bottino C; Moretta A  
 CS Dipartimento di Medicina Sperimentale, Universita di Genova, Genova, Italy.  
 SO HUMAN IMMUNOLOGY, (2000 Nov) 61 (11) 1055-8.  
 Journal code: G9W. ISSN: 0198-8859.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)  
 LA English  
 FS Priority Journals  
 EM 200102  
 ED Entered STN: 20010322  
 Last Updated on STN: 20010322  
 Entered Medline: 20010208  
 AB The lack of classical HLA-class I molecules on trophoblast is necessary to prevent allorecognition by maternal CTL, but may induce activation of NK cells. A protective role against NK cells equipped of suitable inhibitory receptors has been proposed for nonclassical HLA-class I molecules including HLA-E and HLA-G. In the present study we show that the NK-mediated killing of two choriocarcinoma cell lines, JAR and JEG3, is induced upon engagement of natural cytotoxicity receptors (NCR) with their specific ligands. In particular, we show that NKp44, a triggering receptor expressed at the NK cell surface only after in vitro culture in the presence of IL-2, plays a central role in triggering NK cytotoxicity against trophoblast cells. Also NKp46 appear to contribute to this function by cooperating with NKp44. On the other hand, other triggering receptors such as NKp30, 2B4, and **NKG2D** are not involved in killing of choriocarcinoma. Our findings suggest that resistance of trophoblast to NK-mediated cytotoxicity is the result of insufficient activating interactions between the various triggering NK receptors and their target cell ligands. On the other hand, the interaction of nonclassical HLA class I molecules with inhibitory NK receptors appears to play only a marginal role in regulating the susceptibility of choriocarcinoma to NK mediated cytotoxicity.  
 CT Check Tags: Human; Support, Non-U.S. Gov't  
 Antibodies, Monoclonal: IM, immunology

\*Choriocarcinoma: IM, immunology  
 Cytotoxicity Tests, Immunologic  
 \*Cytotoxicity, Immunologic  
 Histocompatibility Antigens Class I: IM, immunology  
 Interleukin-2: PD, pharmacology  
 Killer Cells, Natural: DE, drug effects  
 \*Killer Cells, Natural: IM, immunology  
 \*Receptors, Immunologic: IM, immunology  
 Tumor Cells, Cultured

L8 ANSWER 6 OF 7 MEDLINE  
 AN 2000045048 MEDLINE  
 DN 20045048 PubMed ID: 10574749  
 TI Natural killer cells: stress out, turn on, tune in.  
 AU Diefenbach A; Raulet D H  
 CS Department of Molecular and Cell Biology, Cancer Research Laboratory, 485  
 Life Sciences Addition, University of California at Berkeley, Berkeley,  
 94720-3200, USA.  
 SO CURRENT BIOLOGY, (1999 Nov 18) 9 (22) R851-3. Ref: 14  
 Journal code: B44; 9107782. ISSN: 0960-9822.  
 CY ENGLAND: United Kingdom  
 DT Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LA English  
 FS Priority Journals  
 EM 200005  
 ED Entered STN: 20000518  
 Last Updated on STN: 20000518  
 Entered Medline: 20000511  
 AB Natural killer cells attack **tumor** cells, infected cells and some  
 normal cells, but the basis of their specificity is not completely  
 understood. Recent studies indicate that epithelial **tumor** cells  
 upregulate a stress-induced MHC class-I-like protein termed MICA,  
 triggering NK cells via a recently described receptor called **NKG2D**  
 .  
 CT Check Tags: Animal; Human  
 \*Carcinoma: IM, immunology  
 \*Heat-Shock Proteins: IM, immunology  
 \*Histocompatibility Antigens Class I: IM, immunology  
 \*Killer Cells, Natural: IM, immunology  
 Lymphocyte Transformation  
 Membrane Proteins: CH, chemistry  
 \*Membrane Proteins: IM, immunology  
 Mice  
 \*Models, Immunological  
 \*Neoplasm Proteins: IM, immunology  
 Receptors, Immunologic: CH, chemistry  
 \*Receptors, Immunologic: IM, immunology  
 T-Lymphocytes: IM, immunology

L8 ANSWER 7 OF 7 MEDLINE  
 AN 96235029 MEDLINE  
 DN 96235029 PubMed ID: 8642329  
 TI An autosomal dominant locus, Nka, mapping to the Ly-49 region of a rat  
 natural killer (NK) gene complex, controls NK cell lysis of allogeneic  
 lymphocytes.  
 AU Dissen E; Ryan J C; Seaman W E; Fossum S  
 CS Department of Anatomy, University of Oslo, Norway.  
 SO JOURNAL OF EXPERIMENTAL MEDICINE, (1996 May 1) 183 (5) 2197-207.  
 Journal code: I2V; 2985109R. ISSN: 0022-1007.  
 CY United States  
 DT Journal; Article; (JOURNAL ARTICLE)

LA English  
 FS Priority Journals  
 OS GENBANK-U56822; GENBANK-U56823; GENBANK-U56824; GENBANK-U56863  
 EM 199607  
 ED Entered STN: 19960726  
 Last Updated on STN: 19980206  
 Entered Medline: 19960718

AB Natural Killer (NK) cells can recognize and kill MHC-incompatible normal bone marrow-derived cells. Presently characterized MHC-binding receptors on NK cells, including the Ly-49 family in the mouse, transmit inhibitory signals upon binding to cognate class I MHC ligands. Here we study in vivo NK-mediated lysis of normal allogeneic lymphocytes in crosses between alloreactivity-competent PVG rats and alloreactivity-deficient DA rats. NK cells from both strains are able to lyse standard **tumor** targets. We identify an autosomal dominant locus, Nka, that controls NK-mediated alloreactivity. Individuals carrying the dominant PVG allele in single dose were fully competent in eliminating allogeneic target cells, suggesting that Nka encodes or regulates a gene product inducing or activating alloreactivity. By linkage analysis and pulsed field gel electrophoresis, a natural killer gene complex (NKC) on rat chromosome 4 is described that contains the rat NKR-P1 and Ly-49 multigene families plus a rat NKG2D homologue. Nka maps within the NKC, together with the most telomeric Ly-49 family members, but separate from NKG2D and the NKR-P1 family. The Nka-encoded response, moreover, correlates with the expression of transcripts for Ly-49 receptors in NK cell populations, as Northern blot analysis demonstrated low expression of Ly-49 genes in DA NK cells, in contrast to high expression in alloreactivity-competent PVG, (DA X PVG)F1, and PVG.1AVI NK cells. The low Ly-49 expression in DA is not induced by MHC haplotype, as demonstrated by high expression of Ly-49 in the DA MHC-congenic PVG.1AVI strain. Finally, we have cloned and characterized the first four members of the rat Ly-49 gene family. Their cytoplasmic domains demonstrate substantial heterogeneity, consistent with the hypothesis that different Ly-49 family members may subserve different signaling functions.

CT Check Tags: Animal; Support, Non-U.S. Gov't; Support, U.S. Gov't, Non-P.H.S.; Support, U.S. Gov't, P.H.S.  
 Amino Acid Sequence  
 Antigens, Surface: IM, immunology  
 Base Sequence  
 \*Chromosome Mapping  
 Consensus Sequence  
 Crosses, Genetic  
 DNA Primers  
 Exons  
 \*Genes, Dominant  
 Histocompatibility Antigens Class I: IM, immunology  
 Isoantigens: IM, immunology  
 \*Killer Cells, Natural: IM, immunology  
 Major Histocompatibility Complex  
 Membrane Glycoproteins: BI, biosynthesis  
 \*Membrane Glycoproteins: GE, genetics  
 Mice  
 Molecular Sequence Data  
 Phylogeny  
 Polymerase Chain Reaction  
 Pseudogenes  
 Rats  
 Rats, Inbred F344  
 Rats, Inbred Lew  
 Rats, Inbred Strains  
 Sequence Homology, Amino Acid